arXiv:math/0508055v2 [math.QA] 7 Aug 2005

Multiplier Hopf group coalgebras from algebraic and analytical point of views

A.Hegazi
Mathematics department, Faculty of Science, Mansoura University, Egypt
hegazi@mans.edu.eg
A.Elhafz
Mathematics department, Faculty of Education, Suez Canal University, Egypt
a.elhafz@hotmail.com

Abstract

The Multiplier Hopf Group Coalgebra was introduced by Hegazi in 2002 [7] as a generalization of Hope group caolgebra, introduced by Turaev in 2000 [5], in the non-unital case. We prove that the concepts introduced by A.Van Daele in constructing multiplier Hopf algebra [3] can be adapted to serve again in our construction. A multiplier Hopf group coalgebra is a family of algebras $A = \{A_\alpha\}_{\alpha\in\pi}$, ($\pi$ is a discrete group) equipped with a family of homomorphisms $\Delta = \{\Delta_{\alpha,\beta} : A_{\alpha\beta} \longrightarrow M(A_\alpha \otimes A_\beta)\}_{\alpha,\beta\in\pi}$ which is called a comultiplication under some conditions, where $M(A_\alpha \otimes A_\beta)$ is the multiplier algebra of $A_\alpha \otimes A_\beta$.

In 2003 A. Van Daele suggest a new approach to study the same structure by consider the direct sum of the algebras A_p's which will be a multiplier Hopf algebra called later group cograded multiplier Hope algebra [8]. And hence there exist a one to one correspondence between multiplier Hopf Group Coalgebra and group cograded multiplier Hopf algebra. By using this one-one correspondence we studied multiplier Hopf Group Coalgebra

1 Introduction

In 1992, Alfons Van Daele [3, 4] introduced the notion of multiplier Hopf algebra which is considered a generalization of a usual Hopf algebra. The theory of multiplier Hopf algebra provided us with a mathematical tool for studying noncompact quantum groups with Haar measure.
Recently, Quasitriangular Hopf $\pi-$coalgebras were introduced by Turaev [5]. He showed that

they give rise to crossed $\pi-$categories.Viralizier [6] studied the algebraic properties of the Hopf $\pi-$coalgebras. He showed the existence of integrals and trace for such coalgebras and generalized the main properties of the quasitriangular Hopf algebras to the setting of Hopf $\pi-$coalgebras. Now let us give some basic definitions which we will need in our work: Assume that A is an associative algebra over $\mathbb{C}$. By a left multiplier of A we mean a linear map $L : A \longrightarrow A$ such that $L(ab) = L(a)b$ for all $a, b \in A$. Similarly, by a right multiplier we mean a linear map $R : A \longrightarrow A$ such that $R(ab) = aR(b)$ for all $a, b \in A$. A multiplier (double multiplier) on A is a pair (L, R) such that L is a left multiplier, R is a right multiplier and $R(a)b = aL(b)$ for all $a, b \in A$.

we denoted by $L(A), R(A)$, and $M(A)$ that the set of all left, right, and multipliers of A. It is clear that the composition of maps makes these vector spaces into algebras.

If the product of A is nondegenerate (i.e. if $ab = 0$ for all $a \in A$ then $b = 0$ and if $ab = 0$ for all $b \in A$ then $a = 0$) then the maps $a \longrightarrow \cdot a$, $a \longrightarrow a\cdot$ and $a \longrightarrow (\cdot a, a\cdot)$ give a natural imbedding of A into $L(A), R(A)$ and $M(A)$ respectively.

Since the tensor product of two nondegenerate algebras is again nondegenerate then $A \otimes A \subseteq M(A) \otimes M(A) \subseteq M(A \otimes A)$.

If a homomorphism φ form A into $M(B)$ is nondegenerate (i.e. B is spanned by vectors $\varphi(a)b$ and spanned by vectors $b\varphi(a)$)then it has a unique extension to a homomorphism $M(A) \longrightarrow M(B)$.

- A Hopf algebra over $\mathbb{C}$ (the field of complex numbers) is a tuple $(A, \Delta, \varepsilon, S)$ where A is an associative unital algebra over $\mathbb{C}$, $\Delta : A \longrightarrow A \otimes A$, $\varepsilon : A \longrightarrow \mathbb{C}$ are algebra homomorphisms, and $S : A \longrightarrow A$ is an algebra antihomomorphism such that Δ is coassociative,

  $$(I \otimes \varepsilon)\Delta = (\varepsilon \otimes I)\Delta = I,$$

  and

  $$m(I \otimes S)\Delta = m(S \otimes I)\Delta = \varepsilon 1_A$$

  where m is the multiplication in A. For more details see [1, 2].

- A $\pi-$coalgebra over $\mathbb{C}$ is a family $A = \{A_\alpha\}_{\alpha\in\pi}$($\pi$ is a discrete group) of $\mathbb{C}$ spaces endowed with a family $\Delta = \{\Delta_{\alpha,\beta} : A_{\alpha\beta} \longrightarrow A_\alpha \otimes A_\beta\}$ of $\mathbb{C}-$linear maps (the comultiplication) and a linear map $\varepsilon : A_1 \longrightarrow \mathbb{C}$ (the counit) such that

  1. For any $\alpha, \beta, \gamma \in \pi$; Δ is coassociative in the sense that

     $$(\Delta_{\alpha,\beta} \otimes I)\delta_{\alpha\beta,\gamma} = (I \otimes \Delta_{\beta,\gamma})\Delta_{\alpha,\beta\gamma},$$

  2. For all $\alpha \in \pi$

     $$(I \otimes \varepsilon)\Delta_{\alpha,1} = (\varepsilon \otimes I)\Delta_{1,\alpha} = I$$

     where 1 is the identity in in the group and I is the identity map .

- A Hopf $\pi-$coalgebra over $\mathbb{C}$ is a $\pi-$coalgebra $H = (\{H_\alpha\}_{\alpha\in\pi}, \Delta, \varepsilon)$ endowed with a family $S = \{S_\alpha : H_\alpha \longrightarrow H_{\alpha^{-1}}\}_{\alpha\in\pi}$ of $\mathbb{C}-$linear maps (the antipode)such that

   1. Each H_α is an algebra with multiplication m_α and unit element $1_\alpha \in H_\alpha$,
   2. $\varepsilon : H_1 \longrightarrow \mathbb{C}$ and $\Delta_{\alpha,\beta} : H_{\alpha\beta} \longrightarrow H_\alpha \otimes H_\beta$ (for all $\alpha, \beta \in \pi$) are algebra homomorphisms ,
   3. For any $\alpha \in \pi$,

$$m_\alpha(S_{\alpha^{-1}} \otimes I)\Delta_{\alpha^{-1},\alpha} = \varepsilon 1_\alpha = m_\alpha(I \otimes S_{\alpha^{-1}})\Delta_{\alpha,\alpha^{-1}}.$$

- A multiplier Hopf algebra is a pair (A, Δ) where A is an associative algebra over $\mathbb{C}$, with or without identity and $\Delta : A \longrightarrow M(A \otimes A)$ is an algebra homomorphism, $M(A \otimes A)$ is the multiplier algebra of $A \otimes A$, such that

   1. For all $a, b \in A$

$$\Delta(a)(1 \otimes b) \in A \otimes A \quad \text{and} \quad (a \otimes 1)\Delta(b) \in A \otimes A,$$

   2. Δ is coassociative in the sense that

$$(a \otimes 1 \otimes 1)(\Delta \otimes I)(\Delta(b)(1 \otimes c)) = (I \otimes \Delta)((a \otimes 1)\Delta(b))(1 \otimes 1 \otimes c)$$

   for all $a, b, c, \in A$,

   3. The linear maps

$$\begin{aligned} T^1 &: A \otimes A \longrightarrow A \otimes A \quad , \\ T^2 &: A \otimes A \longrightarrow A \otimes A \end{aligned}$$

   defined by

$$\begin{aligned} T^1(a \otimes b) &= \Delta(a)(1 \otimes b) \quad , \\ T^2(a \otimes b) &= (a \otimes 1)\Delta(b) \end{aligned}$$

   are bijective.

2 Multiplier Hopf group coalgebra.

Assume that $A = \{A_\alpha\}_{\alpha\in\pi}$ (where π is a discrete group) is a family of associative algebras over $\mathbb{C}$ with nondegenerate products, A_α may or may not have a unit.

Definition 2.1. A comultiplication on A is a family of homomorphisms

$$\Delta = \{\Delta_{\alpha,\beta} : A_{\alpha\beta} \longrightarrow M(A_\alpha \otimes A_\beta)\}_{\alpha,\beta\in\pi}$$

such that for any $\alpha, \beta, \gamma \in \pi$

1. $\Delta_{\alpha,\beta}(a)(1 \otimes b) \in A_\alpha \otimes A_\beta \quad \forall a \in A_{\alpha\beta}, b \in A_\beta$ and $(a \otimes 1)\Delta_{\alpha,\beta}(b) \in A_\alpha \otimes A_\beta \quad \forall a \in A_\alpha, b \in A_{\alpha\beta}$,
2. Δ is coassociative in the sense that

   $$(a \otimes 1 \otimes 1)(\Delta_{\alpha,\beta} \otimes I)(\Delta_{\alpha\beta,\gamma}(b)(1 \otimes c)) = (I \otimes \Delta_{\beta,\gamma})((a \otimes 1)\Delta_{\alpha,\beta\gamma}(b))(1 \otimes 1 \otimes c)$$

   for all $a \in A_\alpha,\ b \in A_{\alpha\beta\gamma}$, and $c \in A_\gamma$.

Condition (1) makes sense because $A_\alpha \otimes A_\beta \subseteq M(A_\alpha) \otimes M(A_\beta) \subseteq M(A_\alpha \otimes A_\beta)$. Also, if A_α is unital algebra $\forall \alpha \in \pi$, the above definition gives rise to the definition of comultiplication introduced by Tureav[5].

Definition 2.2. Let $A = \{A_\alpha\}_{\alpha\in\pi}$ be a family of algebras with a nondegenerate products over $\mathbb{C}$, A_α may or may not have a unit and let Δ be a comultiplication on A .We call (A, Δ) a multiplier Hopf $\pi-$coalgebra if the linear maps

$$T^1_{\alpha,\beta} : A_{\alpha\beta} \otimes A_\beta \longrightarrow A_\alpha \otimes A_\beta \quad ,$$
$$T^2_{\alpha,\beta} : A_\alpha \otimes A_{\alpha\beta} \longrightarrow A_\alpha \otimes A_\beta$$

defined by

$$T^1_{\alpha,\beta}(a \otimes b) = \Delta_{\alpha,\beta}(a)(1 \otimes b) \quad ,$$
$$T^2_{\alpha,\beta}(a \otimes b) = (a \otimes 1)\Delta_{\alpha,\beta}(b)$$

are bijective for all $\alpha, \beta \in \ \pi$.

Remark 1. In fact the condition that $T^1_{\alpha,\beta}, T^2_{\alpha,\beta}$ are bijective implies that Δ is nondegenerate homomorphism, and hence the homomorphisms $(\Delta_{\alpha,\beta} \otimes I)$ and $(I \otimes \Delta_{\beta,\gamma})$ have unique extensions to $M(A_{\alpha\beta} \otimes A_\gamma)$ and $M(A_\alpha \otimes A_{\beta\gamma})$ respectively so the coassociativity will just mean

$$(\Delta_{\alpha,\beta} \otimes I)\Delta_{\alpha\beta,\gamma} = (I \otimes \Delta_{\beta,\gamma})\Delta_{\alpha,\beta\gamma}$$

as maps from $A_{\alpha\beta\gamma}$ to $M(A_\alpha \otimes A_\beta \otimes A_\gamma)$.
Also, we can see that $(A_1, \Delta_{1,1})$ is a multiplier Hopf algebra.

Definition 2.3. A multiplier Hopf $\pi-$coalgebra (A, Δ) is called regular multiplier Hopf $\pi-$coalgebra if the opposite comultiplication $\Delta' = \sigma\Delta$ satisfies

1. $\Delta'_{\alpha,\beta}(a)(1\otimes b)\in A_\beta\otimes A_\alpha,\ \ (a\otimes 1)\Delta'_{\alpha,\beta}(b)\in A_\beta\otimes A_\alpha$ for all $\alpha,\beta\in\pi$,
2. Δ' is opposite coassociative in the following sense
$$(a\otimes 1\otimes 1)(\Delta'_{\beta,\alpha}\otimes I)(\Delta'_{\gamma,\beta\alpha}(b)(1\otimes c))=(I\otimes\Delta'_{\gamma,\beta})((a\otimes 1)\Delta'_{\gamma\beta,\alpha}(b))(1\otimes 1\otimes c),$$
3. The linear maps
$$\begin{aligned} T'^{1}_{\alpha,\beta} &: A_{\alpha\beta}\otimes A_\alpha\longrightarrow A_\beta\otimes A_\alpha,\\ T'^{2}_{\alpha,\beta} &: A_\beta\otimes A_{\alpha\beta}\longrightarrow A_\beta\otimes A_\alpha\end{aligned}$$
defined by
$$\begin{aligned} T'^{1}_{\alpha,\beta}(a\otimes b) &= \Delta'_{\alpha,\beta}(a)(1\otimes b),\\ T'^{2}_{\alpha,\beta}(a\otimes b) &= (a\otimes 1)\Delta'_{\alpha,\beta}(b)\end{aligned}$$
are bijective.

Example. Let π be any group, and A be the algebra of complex finitely supported functions on π. In this case $M(A)$ consists of all complex functions on π. Moreover $A\otimes A$ can be naturally identified with finitely supported functions on $\pi\otimes\pi$ so that $M(A\otimes A)$ consists of all complex functions on $\pi\otimes\pi$.
If we define $A_\pi=\{A_\alpha\}_{\alpha\in\pi}$ where $A_\alpha=A$ for all $\alpha\in\pi$,and $\Delta_{\alpha,\beta}:A_{\alpha\beta}\longrightarrow M(A_\alpha\otimes A_\beta)$ where
$$\Delta_{\alpha,\beta}(f)(s,t)=f(\beta^{-1}s\beta t),$$
we will clearly get a family of homomorphisms $\Delta=\{\Delta_{\alpha,\beta}\}_{\alpha,\beta\in\pi}$. If $f\in A_{\alpha\beta},g\in A_\beta$ then $(s,t)\longrightarrow f(\beta^{-1}s\beta t)g(t)$ will have finite support, and if $g\in A_\alpha,f\in A_{\alpha\beta}$ then $(s,t)\longrightarrow g(s)f(s\beta^{-1}t\beta)$ will have finite support. By direct calculations we can see that Δ is coassociative hence it is a comultiplication. The maps $T^1_{\alpha,\beta},T^2_{\alpha,\beta}$ are bijective with inverses $R^1_{\alpha,\beta},R^2_{\alpha,\beta}$ defined by
$$\begin{aligned} R^1_{\alpha,\beta}(f)(s,t) &= f(\beta s t^{-1}\beta^{-1},t),\\ R^1_{\alpha,\beta}(f)(s,t) &= f(s,\beta s^{-1}t\beta^{-1}).\end{aligned}$$

Let us now browse the two approaches for studying the structure of multiplier Hopf group coalgebra.

3 The first approach.

In this section we prove the existence of a suitable counit and antipode for the multiplier Hopf group coalgebra. Also, we prove that there exist a large subspace of the dual space that can be made into an algebra. Finally, we prove that A is a Hopf group coalgebra if and only if it is a unital multiplier Hopf group coalgebra, and that S^{-1} exists under some conditions.

3.1 Construction of the couint.

Definition 3.1. For every $\alpha \in \pi$ define a linear map

$$E_\alpha : A_1 \longrightarrow L(A_\alpha)$$

by

$$E_\alpha(a)b = m_\alpha \left(T^1_{1,\alpha}\right)^{-1}(a \otimes b) \quad \forall \; b \in A_\alpha,$$

where $L(A_\alpha)$ is the left multiplier algebra of A_α, $m_\alpha : A_\alpha \otimes A_\alpha \longrightarrow A_\alpha$ is the multiplication on A_α and $T^1_{1,\alpha}(a \otimes b) = \Delta_{1,\alpha}(a)(1 \otimes b)$.

Because $T^1_{1,\alpha}(x(1 \otimes c)) = T^1_{1,\alpha}(x)(1 \otimes c)$ for all $x \in A_\alpha \otimes A_\alpha, c \in A_\alpha$, the same result holds for $\left(T^1_{1,\alpha}\right)^{-1}$. And since $m_\alpha(x(1 \otimes c)) = m_\alpha(x)c$ we get that $E_\alpha(a)$ is a left multiplier on A_α for all $a \in A_1, \alpha \in \pi$.
Following A.Van Daele we will show that $E_\alpha(A_1) \subset \mathbb{C}1_\alpha$ which will give the map ε.

Lemma 3.2. *For all $a, b \in A_\alpha$, $\alpha \in \pi$ we have*

$$(I \otimes E_\alpha)((b \otimes 1)\Delta_{\alpha,1}(a)) = ba \otimes 1.$$

Proof. Suppose that $a, b \in A_\alpha$, and let

$$a \otimes b = \sum_{i=1}^{n} \Delta_{\alpha,\alpha}(a_i)(1 \otimes b_i).$$

If we apply $(\Delta_{\alpha,1} \otimes I)$ and multiply by $(c \otimes 1 \otimes 1)$ on the left then

$$\begin{aligned}(c \otimes 1)\Delta_{\alpha,1}(a) \otimes b &= \sum (c \otimes 1 \otimes 1)(\Delta_{\alpha,1} \otimes I)(\Delta_{\alpha,\alpha}(a_i)(1 \otimes b_i)) \\ &= \sum (I \otimes \Delta_{1,\alpha})((c \otimes 1)\Delta_{\alpha,\alpha}(a_i))(1 \otimes 1 \otimes b_i).\end{aligned}$$

Now, let φ be any linear functional on A_α. If we apply $(\varphi \otimes I \otimes I)$ to the above equation, we get

$$\begin{aligned}(\varphi \otimes I)((c \otimes 1)\Delta_{\alpha,1}(a)) \otimes b &= (\varphi \otimes I \otimes I)(\sum (I \otimes \Delta_{1,\alpha})((c \otimes 1)\Delta_{\alpha,\alpha}(a_i))(1 \otimes 1 \otimes b_i)) \\ &= \sum \Delta_{1,\alpha}((\varphi \otimes 1)((c \otimes 1)\Delta_{\alpha,\alpha}(a_i)))(1 \otimes b_i) \\ &= \sum T^1_{1,\alpha}((\varphi \otimes I)((c \otimes 1)\Delta_{\alpha,\alpha}(a_i)) \otimes b_i).\end{aligned}$$

By the definition of E_α we have

$$E_\alpha((\varphi \otimes I)((c \otimes 1)\Delta_{\alpha,1}(a)))\, b = \sum (\varphi \otimes I)((c \otimes 1)\Delta_{\alpha,\alpha}(a_i))\, b_i$$

then

$$\begin{aligned}(\varphi \otimes I)((I \otimes E_\alpha)((c \otimes 1)\Delta_{\alpha,1}(a))(1 \otimes b)) &= (\varphi \otimes I)((c \otimes 1) \sum \Delta_{\alpha,\alpha}(a_i)(1 \otimes b_i)) \\ &= (\varphi \otimes I)((c \otimes 1)(a \otimes b)).\end{aligned}$$

Since this holds for all $\varphi \in A_\alpha^{'}$ we get

$$(I \otimes E_\alpha)((c \otimes 1)\Delta_{\alpha,1}(a))(1 \otimes b) = (ca \otimes 1)(1 \otimes b).$$

This gives the required formula in the multiplier algebra.

□

Lemma 3.3. $E_\alpha(A_1) \subset \mathbb{C}1_\alpha \quad \forall \alpha \in \pi$

Proof. By the bijectivity of $T^2_{\alpha,1}$ defined by $T^2_{\alpha,1}(p \otimes q) = (p \otimes 1)\Delta_{\alpha,1}(q)$ we have

$$(I \otimes E_\alpha)(a \otimes b) \in A_\alpha \otimes 1 \ \ \forall \ \ a \in A_1, \ \ b \in A_\alpha.$$

Which gives that $E_\alpha(A_1) \subseteq \mathbb{C}1_\alpha$ for all $\alpha \in \pi$.

□

Definition 3.4. Define for every $\alpha \in \pi$ a map

$$\varepsilon_\alpha : A_1 \longrightarrow \mathbb{C}$$

by

$$\varepsilon_\alpha(a)1_\alpha = E_\alpha(a) \qquad \forall a \in A_1.$$

Now we can rewrite the formula in lemma 3.2 as

$$(I \otimes \varepsilon_\alpha)((a \otimes 1)\Delta_{\alpha,1}(b)) = ab.$$

By the definition of ε_α we have

$$(\varepsilon_\alpha \otimes I)(a \otimes b) = m_\alpha \left(T^1_{1,\alpha}\right)^{-1} (a \otimes b)$$

and hence

$$(\varepsilon_\alpha \otimes I)(\Delta_{1,\alpha}(a)(1 \otimes b) = ab.$$

Lemma 3.5. *ε_α is a homomorphism for all $\alpha \in \pi$*

Proof. Let $a \in A_\alpha$ and $b, c \in A_1$. By the bijectivity of $T^1_{\alpha,1}$ we have

$$\begin{aligned} a \otimes bc &= \sum_i (1 \otimes b)(d_i \otimes 1)\Delta_{\alpha,1}(c_i) \\ &= \sum_{i,j} (a_{ij} \otimes 1)\Delta_{\alpha,1}(b_{ij})\Delta_{\alpha,1}(c_i) \\ &= \sum_{i,j} (a_{ij} \otimes 1)\Delta_{\alpha,1}(b_{ij}c_i) \end{aligned}$$

where $a_{ij}, b_{ij}, c_i, d_i \in A_\alpha$. Applying $(I \otimes \varepsilon_\alpha)$ we get

$$a\varepsilon_\alpha(bc) = \sum_{i,j} a_{ij}b_{ij}c_i.$$

On the other hand we have

$$a\varepsilon_\alpha(c) = \sum_i d_i c_i$$

if

$$a \otimes c = \sum_i (d_i \otimes 1)\Delta_{\alpha,1}(c_i).$$

Then

$$\begin{aligned} a \otimes b\varepsilon_\alpha(c) &= \sum_i (d_i \otimes b)(c_i \otimes 1) \\ &= \sum_{i,j} (a_{ij} \otimes 1)\Delta_{\alpha,1}(b_{ij})(c_i \otimes 1). \end{aligned}$$

Applying $(I \otimes \varepsilon_\alpha)$ we get

$$a\varepsilon_\alpha(b)\varepsilon_\alpha(c) = \sum_{i,j} a_{ij}b_{ij}c_i.$$

This means

$$a\varepsilon_\alpha(bc) = a\varepsilon_\alpha(b)\varepsilon_\alpha(c).$$

□

Lemma 3.6. *For all* $\alpha, \beta \in \pi \qquad \varepsilon_\alpha = \varepsilon_\beta$.

Proof. We can see that the map

$$\varphi : E_\alpha(A_1) \longrightarrow E_\beta(A_1)$$

defined by

$$\varphi(E_\alpha(a)) = E_\beta(a) \quad \forall a \in A_1$$

is an algebra homomorphism. Then

$$\varepsilon_\beta(a)I_\beta = E_\beta(a) = \varphi(E_\alpha(a)) = \varphi(\varepsilon_\alpha(a)I_\alpha) = \varepsilon_\alpha(a)\varphi(I_\alpha) = \varepsilon_\alpha(a)I_\beta.$$

Since this holds for all $a \in A_1$, then $\varepsilon_\alpha = \varepsilon_\beta$ for all $\alpha, \beta \in \pi$. □

Now, we can summarize the results in this section in the following theorem(generalize [3] theorem 3.6).

Theorem 1. *Let* (A, Δ) *be a multiplier Hopf* $\pi-$*coalgebra. Then there exist a homomorphism* $\varepsilon : A_1 \longrightarrow C$ *such that*

$$(I \otimes \varepsilon)((a \otimes 1)\Delta_{\alpha,1}(b)) = ab,$$

$$(\varepsilon \otimes I)(\Delta_{1,\alpha}(a)(1 \otimes b)) = ab$$

for all $a, b \in A_\alpha,\ \alpha \in \pi$.

Back to our example when $f \in A_1, g \in A_\alpha$ and $t \in \pi$

$$\varepsilon(f)g(t) = (m_\alpha(T_{1,\alpha})^{-1}(f \otimes g))(t) = (f \otimes g)(\alpha t t^{-1}\alpha^{-1}, t) = f(e)g(t)$$

and hence $\varepsilon(f) = f(e)$.

3.2 Construction of the antipode.

Definition 3.7. For every $\alpha \in \pi$ define a map

$$S_\alpha : A_\alpha \longrightarrow L(A_{\alpha^{-1}})$$

by

$$S_\alpha(a)b = (\varepsilon_\alpha \otimes I)(T^1_{\alpha,\alpha^{-1}})^{-1}(a \otimes b) \quad \forall \ b \in A_{\alpha^{-1}}.$$

As before $S_\alpha(a)$ is a left multiplier on $A_{\alpha^{-1}}$ for all $a \in A_\alpha$.

Lemma 3.8. *For all* $a \in A_1,\ b, c \in A_\alpha,\ \alpha \in \pi$, *we have*

$$(I \otimes S_{\alpha^{-1}})((c \otimes 1)\Delta_{\alpha,\alpha^{-1}}(a))(1 \otimes b) = (c \otimes 1)(T^1_{1,\alpha})^{-1}(a \otimes b).$$

Proof. Assume that $a \in A_1,\ b \in A_\alpha$. Write

$$a \otimes b = \sum_{i=1}^{n} \Delta_{\alpha\alpha^{-1},\alpha}(a_i)(1 \otimes b_i).$$

Applying $(\Delta_{\alpha,\alpha^{-1}} \otimes I)$, and multiplying by $(c \otimes 1 \otimes 1)$ on the left, we obtain

$$\begin{aligned} (c \otimes 1)\Delta_{\alpha,\alpha^{-1}}(a) \otimes b &= \sum (c \otimes 1 \otimes 1)(\Delta_{\alpha,\alpha^{-1}} \otimes I)(\Delta_{\alpha\alpha^{-1},\alpha}(a_i)(1 \otimes b_i)) \\ &= \sum (I \otimes \Delta_{\alpha^{-1},\alpha})((c \otimes 1)\Delta_{\alpha,\alpha^{-1}\alpha}(a_i))(1 \otimes 1 \otimes b_i). \end{aligned}$$

Let $\varphi \in A'_\alpha$, and apply $(\varphi \otimes I \otimes I)$ then

$$\begin{aligned} (\varphi \otimes I)((c \otimes 1)\Delta_{\alpha,\alpha^{-1}}(a)) \otimes b &= \sum \Delta_{\alpha^{-1},\alpha}((\varphi \otimes I)((c \otimes 1)\Delta_{\alpha,\alpha^{-1}\alpha}(a_i)))(1 \otimes b_i) \\ &= \sum T^1_{\alpha^{-1},\alpha}((\varphi \otimes I)((c \otimes 1)\Delta_{\alpha,\alpha^{-1}\alpha}(a_i)) \otimes b_i). \end{aligned}$$

By the definition of $S_{\alpha^{-1}}$ we have

$$\begin{aligned} S_{\alpha^{-1}}((\varphi\otimes I)((c\otimes 1)\Delta_{\alpha,\alpha^{-1}}(a))b &= \sum(\varepsilon\otimes I)((\varphi\otimes I)((c\otimes 1)\Delta_{\alpha,\alpha^{-1}\alpha}(a_i))\otimes b_i) \\ &= \sum(\varphi\otimes I)((I\otimes\varepsilon)((c\otimes 1)\Delta_{\alpha,\alpha^{-1}\alpha}(a_i))\otimes b_i) \\ &= (\varphi\otimes I)(\sum ca_i\otimes b_i), \end{aligned}$$

and hence

$$(\varphi\otimes I)((I\otimes S_{\alpha^{-1}})((c\otimes 1)\Delta_{\alpha,\alpha^{-1}}(a))(1\otimes b)) = (\varphi\otimes I)((c\otimes 1)(T^1_{1,\alpha})^{-1}(a\otimes b)).$$

Since this is true for all $\varphi\in A'_\alpha$, we get

$$(I\otimes S_{\alpha^{-1}})((c\otimes 1)\Delta_{\alpha,\alpha^{-1}}(a))(1\otimes b) = (c\otimes 1)(T^1_{1,\alpha})^{-1}(a\otimes b).$$

□

Lemma 3.9. *For all* $a\in A_1,\ \ b,c\in A_\alpha,\ \ \alpha\in\pi,$ *we have*

$$m_\alpha(I\otimes S_{\alpha^{-1}})((c\otimes 1)\Delta_{\alpha,\alpha^{-1}}(a))(1\otimes b) = c\varepsilon(a)b.$$

Proof. If we apply m_α to the equation in lemma 3.8, we get

$$\begin{aligned} m_\alpha((I\otimes S_{\alpha^{-1}})((c\otimes 1)\Delta_{\alpha,\alpha^{-1}}(a))(1\otimes b)) &= m_\alpha((c\otimes 1)(T^1_{1,\alpha})^{-1}(a\otimes b)) \\ &= c\varepsilon(a)b. \end{aligned}$$

□

Lemma 3.10. $S_\alpha(a)$ *is a right multiplier for all* $a\in A_\alpha,\ \ \alpha\in\pi,$ *and satisfies*

$$m_\alpha((c\otimes 1)(S_{\alpha^{-1}}\otimes I)\Delta_{\alpha^{-1},\alpha}(a)(1\otimes b)) = c\varepsilon(a)b$$

for all $a\in A_1,\ \ b,c\in A_\alpha,\ \ \alpha\in\pi.$

Proof. For every $\alpha\in\pi$ define

$$S'_\alpha : A_\alpha \longrightarrow R(A_{\alpha^{-1}})\quad\forall\ \ b\in A_{\alpha^{-1}}$$

by

$$bS'_\alpha(a) = (I\otimes\varepsilon)(T^2_{\alpha^{-1},\alpha})^{-1}(b\otimes a)$$

where $R(A_\alpha)$ is the right multiplier algebra. Directly we can show that $S'_\alpha(a)$ is a right multiplier on $A_{\alpha^{-1}}$ for all $a\in A_\alpha$. Completely similar as in lemma 3.8 we can prove that

$$(c\otimes 1)(S'_{\alpha^{-1}}\otimes I)(\Delta_{\alpha^{-1},\alpha}(a)(1\otimes b)) = (T^2_{\alpha,1})^{-1}(c\otimes a)(1\otimes b).$$

If we apply m_α to the above equation, we get the formula in the statement of the lemma with $S^{'}$ instead of S since

$$m_\alpha((T^2_{\alpha,1})^{-1}(c\otimes a)) = c\varepsilon(a).$$

Now, we will show that $S^{'}_\alpha = S_\alpha$ for all $\alpha\in\pi$.
By definition of $S^{'}$ we have

$$aS^{'}_{\alpha^{-1}}(b) = \sum a_i\varepsilon(b_i)$$

if

$$a\otimes b = \sum (a_i\otimes 1)\Delta_{\alpha,\alpha^{-1}}(b_i).$$

Applying $(I\otimes S_{\alpha^{-1}})$, and multiplying by $1\otimes c$ we get

$$a\otimes S_{\alpha^{-1}}(b)c = \sum (I\otimes S_{\alpha^{-1}})((a_i\otimes 1)\Delta_{\alpha,\alpha^{-1}}(b_i))(1\otimes c).$$

Applying m_α we have

$$aS_{\alpha^{-1}}(b)c = \sum a_i\varepsilon(b_i)c = aS^{'}_{\alpha^{-1}}(b)c,$$

which shows that $S^{'}_\alpha = S_\alpha$for all $\alpha\in\pi$, thus $S_\alpha(a)$ is a multiplier on $A_{\alpha^{-1}}$. ☐

Lemma 3.11. *For all $a, b\in A_\alpha,\ \ \alpha\in\pi$*

$$S_\alpha(ab) = S_\alpha(b)S_\alpha(a).$$

Proof. Let $a, b\in A_{\alpha^{-1}}$, $c\in A_\alpha$. Then

$$\begin{aligned} c\otimes ab &= \sum_i (e_i\otimes a)\Delta_{\alpha,\alpha^{-1}}(b_i) \\ &= \sum_{i,j} (c_{ij}\otimes 1)\Delta_{\alpha,\alpha^{-1}}(a_{i,j})\Delta_{\alpha,\alpha^{-1}}(b_i). \end{aligned}$$

Applying $(I\otimes S_{\alpha^{-1}})$, multiplying by $(1\otimes d)$, and applying m_α we get

$$\begin{aligned} cS_{\alpha^{-1}}(ab)d &= \sum_{i,j} m_\alpha((I\otimes S_{\alpha^{-1}})((c_{ij}\otimes 1)\Delta_{\alpha,\alpha^{-1}}(a_{i,j}))(1\otimes d)) \\ &= \sum_{i,j} c_{ij}\varepsilon(a_{ij}b_i)d. \end{aligned}$$

On the other hand, we have

$$cS_{\alpha^{-1}}(b) = \sum_i e_i\varepsilon(b_i)$$

if

$$c\otimes b = \sum_i (e_i\otimes 1)\Delta_{\alpha,\alpha^{-1}}(b_i).$$

Then

$$\begin{aligned} cS_{\alpha^{-1}}(b) \otimes a &= \sum_i e_i \otimes a\varepsilon(b_i) \\ &= \sum_{i,j} (c_{ij}\varepsilon(b_i) \otimes 1)\Delta_{\alpha,\alpha^{-1}}(a_{i,j}). \end{aligned}$$

Applying $(I \otimes S_{\alpha^{-1}})$, multiplying by $(1 \otimes d)$, and applying m_α we get

$$cS_{\alpha^{-1}}(b)S_{\alpha^{-1}}(a)d = \sum_{i,j} c_{ij}\varepsilon(a_{ij})\varepsilon(b_i)d.$$

Thus

$$S_{\alpha^{-1}}(ab) = S_{\alpha^{-1}}(b)S_{\alpha^{-1}}(a)$$

for all $a, b \in A_{\alpha^{-1}}$ and hence S is an antihomomorphism. $\square$

The following theorem gives the main result of this section [generalize [3] theorem 4.6].

Theorem 2. *If (A, Δ) is a multiplier Hopf $\pi-$coalgebra then there exist an antihomomorphism $S = \{S_\alpha : A_\alpha \longrightarrow M(A_{\alpha^{-1}})\}_{\alpha\in\pi}$ such that for all $a \in A_1 \quad b, c \in A_\alpha \quad \alpha \in \pi$*

$$m_\alpha((I \otimes S_{\alpha^{-1}})((c \otimes 1)\Delta_{\alpha,\alpha^{-1}}(a))(1 \otimes b)) = c\varepsilon(a)b,$$

$$m_\alpha((c \otimes 1)(S_{\alpha^{-1}} \otimes I)\Delta_{\alpha^{-1},\alpha}(a)(1 \otimes b)) = c\varepsilon(a)b.$$

Back to our example let $f \in A_\alpha, g \in A_{\alpha^{-1}}$, and $t \in \pi$. Then

$$\begin{aligned} (S_\alpha(f)g)(t) &= (\varepsilon \otimes I)(T^1_{\alpha,\alpha^{-1}})^{-1}(f \otimes g)(t) = (T^1_{\alpha,\alpha^{-1}})^{-1}(f \otimes g)(e, t) \\ &= (f \otimes g)(\alpha^{-1}t^{-1}\alpha, t) = f(\alpha^{-1}t^{-1}\alpha)g(t) \end{aligned}$$

and hence $S_\alpha(f)(t) = f(\alpha^{-1}t^{-1}\alpha)$.

Theorem 3. *A is a Hopf $\pi-$coalgebra if and only if A is a multiplier Hopf $\pi-$coalgebra with unital component.*

Proof. If (A, Δ) is a Hopf $\pi-$coalgebra then A_α has an identity $\forall\alpha \in \pi$ which means that $M(A_\alpha) = A_\alpha \quad \forall\alpha \in \pi$ and according to remark (1) we have Δ is a comultiplication.
Now we will prove that the linear maps

$$\begin{aligned} T^1_{\alpha,\beta} &: A_{\alpha\beta} \otimes A_\beta \longrightarrow A_\alpha \otimes A_\beta \quad , \\ T^2_{\alpha,\beta} &: A_\alpha \otimes A_{\alpha\beta} \longrightarrow A_\alpha \otimes A_\beta \end{aligned}$$

defined by

$$\begin{aligned} T^1_{\alpha,\beta}(a \otimes b) &= \Delta_{\alpha,\beta}(a)(1 \otimes b) \quad , \\ T^2_{\alpha,\beta}(a \otimes b) &= (a \otimes 1)\Delta_{\alpha,\beta}(b) \end{aligned}$$

are bijective for all $\alpha, \beta \in \pi$.
Consider the following linear maps

$$\begin{aligned} R^1_{\alpha,\beta} : A_\alpha \otimes A_\beta &\longrightarrow A_{\alpha\beta} \otimes A_\beta \quad , \\ R^2_{\alpha,\beta} : A_\alpha \otimes A_\beta &\longrightarrow A_\alpha \otimes A_{\alpha\beta} \end{aligned}$$

which defined by

$$\begin{aligned} R^1_{\alpha,\beta}(a \otimes b) &= ((I \otimes S_{\beta^{-1}})\Delta_{\alpha\beta,\beta^{-1}}(a))(1 \otimes b) \quad , \\ R^2_{\alpha,\beta}(a \otimes b) &= (a \otimes 1)((S_{\alpha^{-1}} \otimes I)\Delta_{\alpha^{-1},\alpha\beta}(b)) \quad . \end{aligned}$$

By using the properties of S and Δ we can prove that $R^1_{\alpha,\beta}$ and $R^2_{\alpha,\beta}$ are the inverses of $T^1_{\alpha,\beta}$ and $T^2_{\alpha,\beta}$ respectively. If we use Sweedler's notation we get

$$\begin{aligned} T^1_{\alpha,\beta}R^1_{\alpha,\beta}(a \otimes b) &= \sum_{(a)} T^1_{\alpha,\beta}(a_{(1,\alpha\beta)} \otimes S_{\beta^{-1}}(a_{(2,\beta^{-1})})\, b) \\ &= \sum_{(a)} a_{(1,\alpha)} \otimes a_{(2,\beta)} S_{\beta^{-1}}(a_{(3,\beta^{-1})})\, b \\ &= \sum_{(a)} a_{(1,\alpha)} \otimes m_\alpha(I \otimes S_{\beta^{-1}})\Delta_{\beta,\beta^{-1}}(a_{(2,1)})(1 \otimes b) \\ &= \sum_{(a)} (a_{(1,\alpha)} \otimes \varepsilon(a_{(2,1)})1_\beta)(1 \otimes b) \;=\; a \otimes b. \end{aligned}$$

Similarly, for $R^1_{\alpha,\beta}T^1_{\alpha,\beta}$, $R^2_{\alpha,\beta}T^2_{\alpha,\beta}$ and $T^2_{\alpha,\beta}R^2_{\alpha,\beta}$. Therefore A is a multiplier Hopf $\pi-$coalgebra.
If A is a unital multiplier Hopf $\pi-$coalgebra then $M(A_\alpha \otimes A_\beta) = A_\alpha \otimes A_\beta \;\forall\, \alpha, \beta \in \pi$ and by using remark (1) we have

$$(\Delta_{\alpha,\beta} \otimes I)\Delta_{\alpha\beta,\gamma} = (I \otimes \Delta_{\beta,\gamma})\Delta_{\alpha,\beta\gamma}$$

as a maps from $A_{\alpha\beta\gamma}$ into $A_\alpha \otimes A_\beta \otimes A_\gamma$
Also, since ε is a nondegenerate homomorphism then $\varepsilon \otimes I$ and $I \otimes \varepsilon$ have unique extensions to $M(A_1 \otimes A_\alpha)$ and $M(A_\alpha \otimes A_1)$, respectively. So the counitary property just means

$$(I \otimes \varepsilon)\Delta_{\alpha,1} = (\varepsilon \otimes 1)\Delta_{1,\alpha} = I \quad \forall \alpha \in \pi.$$

Since A is unital then $M(A_1 \otimes A_\alpha) = A_1 \otimes A_\alpha$ and $M(A_\alpha \otimes A_1) = A_\alpha \otimes A_1$ and then we obtain the definition of the counit introduced by Turaev.
If A is unital then the antihomomorphism S has the form

$$S = \{S_\alpha : A_\alpha \longrightarrow A_{\alpha^{-1}}\}_{\alpha \in \pi}$$

such that

$$m_\alpha(S_{\alpha^{-1}} \otimes I)\Delta_{\alpha^{-1},\alpha} = \varepsilon 1_\alpha = m_\alpha(I \otimes S_{\alpha^{-1}})\Delta_{\alpha,\alpha^{-1}}.$$

□

3.3 Regular multiplier Hopf group coalgebra.

In this subsection, we reformulate the concept of regular multiplier Hopf algebra to obtain a bijective antipode in our structure using the technique of proofs of A. Van Daele.

Definition 3.12. A multiplier Hopf $\pi-$coalgebra (A, Δ) is called regular multiplier Hopf $\pi-$coalgebra if the opposite comultiplication $\Delta^{'} = \sigma\Delta$ satisfies

1. $\Delta^{'}_{\alpha,\beta}(a)(1 \otimes b) \in A_\beta \otimes A_\alpha$, $(a \otimes 1)\Delta^{'}_{\alpha,\beta}(b) \in A_\beta \otimes A_\alpha$ for all $\alpha, \beta \in \pi$,
2. $\Delta^{'}$ is opposite coassociative in the following sense
$$(a \otimes 1 \otimes 1)(\Delta^{'}_{\beta,\alpha} \otimes I)(\Delta^{'}_{\gamma,\beta\alpha}(b)(1 \otimes c)) = (I \otimes \Delta^{'}_{\gamma,\beta})((a \otimes 1)\Delta^{'}_{\gamma\beta,\alpha}(b))(1 \otimes 1 \otimes c),$$
3. The linear maps
$$\begin{aligned} T^{'1}_{\alpha,\beta} &: A_{\alpha\beta} \otimes A_\alpha \longrightarrow A_\beta \otimes A_\alpha, \\ T^{'2}_{\alpha,\beta} &: A_\beta \otimes A_{\alpha\beta} \longrightarrow A_\beta \otimes A_\alpha \end{aligned}$$
defined by
$$\begin{aligned} T^{'1}_{\alpha,\beta}(a \otimes b) &= \Delta^{'}_{\alpha,\beta}(a)(1 \otimes b), \\ T^{'2}_{\alpha,\beta}(a \otimes b) &= (a \otimes 1)\Delta^{'}_{\alpha,\beta}(b) \end{aligned}$$
are bijective.

Lemma 3.13. *For all $\alpha \in \pi$ and $a, b \in A_\alpha$ we have*

- $(\varepsilon \otimes I)((1 \otimes a)\Delta_{1,\alpha}(b)) = ab,$
- $(I \otimes \varepsilon)(\Delta_{\alpha,1}(a)(b \otimes 1)) = ab.$

Proof. Define
$$E^{'}_\alpha : A_1 \to L(A_\alpha)$$
by
$$E^{'}_\alpha(a)b = m_\alpha(T^{'1}_{\alpha,1})^{-1}(a \otimes b).$$
Analogous to lemma 3.2 and using the opposite coassociativity we can prove that
$$(I \otimes E^{'}_\alpha)((c \otimes 1)\Delta^{'}_{1,\alpha}(a)) = ca \otimes 1.$$
Define
$$\varepsilon^{'}_\alpha(a)1_\alpha = E^{'}_\alpha(a).$$

Then

$$(I \otimes \varepsilon^{'})((c \otimes 1)\Delta^{'}_{1,\alpha}(a_i)) = ca.$$

By the definition of $\varepsilon^{'}_{\alpha}$ we have

$$(\varepsilon^{'}_{\alpha} \otimes I)(a \otimes b) = m_{\alpha}(T^{'1}_{\alpha,1})^{-1}(a \otimes b)$$

and hence

$$(\varepsilon^{'}_{\alpha} \otimes I)(\Delta^{'}_{\alpha,1}(a)(b \otimes 1)) = ab.$$

If we apply σ to the above equations we get

$$\begin{aligned}(\varepsilon^{'}_{\alpha} \otimes I)((1 \otimes a)\Delta_{1,\alpha}(b)) &= ab,\\ (I \otimes \varepsilon^{'}_{\alpha})(\Delta_{\alpha,1}(a)(b \otimes 1)) &= ab.\end{aligned}$$

which are the equations in the statement of the lemma with $\varepsilon^{'}_{\alpha}$ instead of ε. We will now prove that $\varepsilon^{'}_{\alpha} = \varepsilon$.
By theorem 1 we have

$$c(\varepsilon_{\alpha} \otimes I)(\Delta_{1,\alpha}(a)(1 \otimes b)) = cab = (\varepsilon_{\alpha} \otimes I)((1 \otimes c)\Delta_{1,\alpha}(a))b.$$

Then

$$(\varepsilon_{\alpha} \otimes I)((1 \otimes c)\Delta_{1,\alpha}(a)) = (\varepsilon^{'}_{\alpha} \otimes I)((1 \otimes c)\Delta_{1,\alpha}(a)).$$

By the bijectivity of the map $\sigma T^{'2}_{\alpha,\beta}$ we have

$$\varepsilon_{\alpha} = \varepsilon^{'}_{\alpha} \qquad \forall\ \alpha \in \pi.$$

□

Definition 3.14. For every $\alpha \in \pi$ define a linear map

$$S^{'}_{\alpha}(a) : A_{\alpha} \to M(A_{\alpha^{-1}})$$

by

$$S^{'}_{\alpha}(a)(b) = (\varepsilon \otimes I\ )(T^{'1}_{\alpha^{-1},\alpha})^{-1}(a \otimes b).$$

We define $S^{'} = \{S^{'}_{\alpha}\}_{\alpha\in\pi}$.

We will show that $S^{'}$ has the same properties of S in the sense of the definition of the regular multiplier Hopf group coalgebra and is the inverse of S.

Lemma 3.15. *If (A, Δ) is a regular multiplier Hopf group coalgebra then S is invertible with inverse $S^{'}$, $S_{\alpha}(A_{\alpha}) \subseteq A_{\alpha^{-1}}$, and $S^{'}_{\alpha}(A_{\alpha}) \subseteq A_{\alpha^{-1}}$ for all $\alpha \in \pi$.*

Proof. Analogous to lemmas 3.8, 3.9, and 3.10; and using the opposite coassociativity we can proof

$$m_\alpha((I\otimes S'_{\alpha^{-1}})((c\otimes 1)\Delta_{\alpha^{-1},\alpha}(a))(1\otimes b)) = c\varepsilon(a)b.$$

Similarly, we can prove that $S'_{\alpha^{-1}}(a)$ is a right multiplier for all $a\in A_\alpha$, $\alpha\in\pi$ and satisfies

$$m_\alpha((c\otimes 1)(S'_{\alpha^{-1}}\otimes I))((\Delta'_{\alpha,\alpha^{-1}}(a)(1\otimes b))) = c\varepsilon(a)b.$$

Now by the definition of S_α we have

$$S_\alpha(a)(b) = \sum \varepsilon_\alpha(a_i)b_i$$

if

$$a\otimes b = \sum \Delta_{\alpha,\alpha^{-1}}(a_i)(1\otimes b_i).$$

Then

$$\begin{aligned} b\otimes a &= \sum \Delta'_{\alpha^{-1},\alpha}(a_i)(b_i\otimes 1),\\ b\otimes ac &= \sum \Delta'_{\alpha^{-1},\alpha}(a_i)(b_i\otimes c). \end{aligned}$$

Applying $(S_{\alpha^{-1}}\otimes I)$ and multiplying by $(d\otimes 1)$

$$\begin{aligned} dS_{\alpha^{-1}}(b)\otimes ac &= \sum (d\otimes 1)(S'_{\alpha^{-1}}\otimes I)(\Delta'_{\alpha,\alpha^{-1}}(a_i)(b_i\otimes c))\\ &= \sum (d\otimes 1)(S'_{\alpha^{-1}}(b_i)\otimes 1)(S'_{\alpha^{-1}}\otimes I)(\Delta'_{\alpha,\alpha^{-1}}(a_i)(1\otimes c)). \end{aligned}$$

Applying m_α

$$\begin{aligned} dS_{\alpha^{-1}}(b)ac &= \sum dS'_{\alpha^{-1}}(b_i)\varepsilon(a_i)c\\ &= \sum dS'_{\alpha^{-1}}(\varepsilon(a_i)b_i)c\\ &= dS'_{\alpha^{-1}}(S_\alpha(a)b)c. \end{aligned}$$

Which shows that

$$S'_{\alpha^{-1}}(b)a = S'_{\alpha^{-1}}(S_\alpha(a)b).$$

Since the elements of the form $S_\alpha(a)b$ generates $A_{\alpha^{-1}}$ then the above formula implies that $S'_{\alpha^{-1}}(A_{\alpha^{-1}})\subseteq A_\alpha$ for all $\alpha\in\pi$. Similarly, $S_{\alpha^{-1}}(A_{\alpha^{-1}})\subseteq A_\alpha$. The above formula gives that

$$S'_{\alpha^{-1}}(b)a = S'_{\alpha^{-1}}(b)S'_{\alpha^{-1}}(S_\alpha(a)).$$

If we multiply by c on the left and use that elements of the form $cS'_{\alpha^{-1}}(b)$ span A, we get $a = S'_{\alpha^{-1}}(S_\alpha(a))$. Similarly, we can show that $b = S_\alpha(S'_{\alpha^{-1}}(b))$. $\square$

Lemma 3.16. *1. Let $a \in A_{\alpha\beta}$, $b \in A_{\beta}$, $a_i \in A_{\alpha}$, and $b_i \in A_{\beta^{-1}}$. Then*

$$a \otimes S_{\beta}(b) = \sum \Delta_{\alpha\beta,\beta^{-1}}(a_i)(1 \otimes b_i) \Leftrightarrow \sum a_i \otimes S_{\beta}^{-1}(b_i) = (1 \otimes b)\Delta_{\alpha,\beta}(a).$$

2. Let $a \in A_{\alpha\beta}$, $b \in A_{\alpha}$, $a_i \in A_{\beta}$, and $b_i \in A_{\alpha^{-1}}$. Then

$$S_{\alpha^{-1}}(b) \otimes a = \sum (b_i \otimes 1)\Delta_{\alpha^{-1},\alpha\beta}(a_i) \Leftrightarrow \Delta_{\alpha,\beta}(a)(b \otimes 1) = \sum S_{\alpha}^{-1}(b_i) \otimes a_i.$$

Proof. Assume that

$$a \otimes S_{\beta}(b) = \sum \Delta_{\alpha\beta,\beta^{-1}}(a_i)(1 \otimes b_i).$$

Applying $(\Delta_{\alpha,\beta} \otimes I)$ and multiplying by $(c \otimes 1 \otimes 1)$ on the left we get

$$\begin{aligned} (c \otimes 1)\Delta_{\alpha,\beta}(a) \otimes S_{\beta}(b) &= \sum (c \otimes 1 \otimes 1)(\Delta_{\alpha,\beta} \otimes I)(\Delta_{\alpha\beta,\beta^{-1}}(a_i)(1 \otimes b_i)) \\ &= \sum (I \otimes \Delta_{\beta,\beta^{-1}})((c \otimes 1)\Delta_{\alpha,1}(a_i))(1 \otimes 1 \otimes b_i). \end{aligned}$$

Let $\varphi \in A_{\alpha}^{'}$. Apply $(\varphi \otimes I \otimes I)$, we get

$$\begin{aligned} (\varphi \otimes I)((c \otimes 1)\Delta_{\alpha,\beta}(a)) \otimes S_{\beta}(b) &= \sum \Delta_{\beta,\beta^{-1}}((\varphi \otimes I)((c \otimes 1)\Delta_{\alpha,1}(a_i)))(1 \otimes b_i) \\ &= \sum T_{\beta,\beta^{-1}}^{'1}((\varphi \otimes I)((c \otimes 1)\Delta_{\alpha,1}(a_i)) \otimes b_i). \end{aligned}$$

By the definition of S_{β} we have

$$\begin{aligned} S_{\beta}((\varphi \otimes I)((c \otimes 1)\Delta_{\alpha,\beta}(a)))S_{\beta}(b)) &= \sum \varepsilon((\varphi \otimes I)((c \otimes 1)\Delta_{\alpha,1}(a_i))) \otimes b_i \\ &= \sum (\varphi \otimes I)((I \otimes \varepsilon)((c \otimes 1)\Delta_{\alpha,1}(a_i)) \otimes b_i). \end{aligned}$$

Then

$$(\varphi \otimes I)((I \otimes S_{\beta})((c \otimes 1)\Delta_{\alpha,\beta}(a))(1 \otimes S_{\beta}(b))) = \sum (\varphi \otimes I)(ca_i \otimes b_i).$$

Since this holds for all $\varphi \in A_{\alpha}'$ we get

$$(I \otimes S_{\beta})(c \otimes b)\Delta_{\alpha,\beta}(a) = \sum (ca_i \otimes b_i).$$

By cancelling c we get

$$(I \otimes S_{\beta})(1 \otimes b)\Delta_{\alpha,\beta}(a) = \sum (a_i \otimes b_i).$$

Applying $I \otimes S_{\beta}^{-1}$ we have

$$(1 \otimes b)\Delta_{\alpha,\beta}(a) = \sum a_i \otimes S_{\beta}^{-1}(b_i).$$

Similarly, we can prove part (2) . ☐

Lemma 3.17. *Let $a \in A_{\alpha^{-1}}$, $b \in A_{\beta^{-1}\alpha^{-1}}$, $a_i \in A_{\alpha^{-1}}$ and $b_i \in A_{\beta}$. Then the following statements are equivalent :*

1. $\Delta_{\beta,\beta^{-1}\alpha^{-1}}(a)(1\otimes b)=\sum\Delta_{\beta,\beta^{-1}\alpha^{-1}}(a_i)(b_i\otimes 1)$,

2. $a\otimes S^{-1}_{\alpha\beta}(b)=\sum(a_i\otimes 1)\Delta_{\alpha^{-1},\alpha\beta}(b_i)$.

Proof. Let

$$\Delta_{\beta,\beta^{-1}\alpha^{-1}}(a)(b\otimes 1)=\sum\Delta_{\beta,\beta^{-1}\alpha^{-1}}(a_i)(b_i\otimes 1).$$

For $c\in A_{\alpha\beta}$ write

$$b_i\otimes S_{\alpha\beta}(c)=\sum_k\Delta_{\beta,\beta^{-1}\alpha^{-1}}(p_{ik})(1\otimes q_{ik})$$

for all i. Then

$$\Delta_{\beta,\beta^{-1}\alpha^{-1}}(a)(1\otimes bS_{\alpha\beta}(c))=\sum_{i,k}\Delta_{\beta,\beta^{-1}\alpha^{-1}}(a_ip_{ik})(1\otimes q_{ik})$$

and by the bijectivity of the map $T^1_{\beta,\beta^{-1}\alpha^{-1}}$ we have

$$a\otimes bS_{\alpha\beta}(c)=\sum_{i,k}a_ip_{ik}\otimes q_{ik}.$$

On the other hand, by Lemma 3.15 we have

$$(1\otimes c)\Delta_{\alpha^{-1},\alpha\beta}(b)=\sum_{i,k}p_{ik}\otimes S^{-1}_{\alpha\beta}(q_{ik}).$$

Therefore

$$\begin{aligned}a\otimes cS^{-1}_{\alpha\beta}(b)&=\sum_{i,k}a_ip_{ik}\otimes S^{-1}_{\alpha\beta}(q_{ik})\\&=\sum_i(a_i\otimes c)\Delta_{\alpha^{-1},\alpha\beta}(b_i).\end{aligned}$$

By cancelling c we get the required formula. ☐

Analogous to lemma 4.9 we can prove the following lemma.

Lemma 3.18. *1. Let $a\in A_{\beta^{-1}\alpha^{-1}}$, $b\in A_{\beta^{-1}}$, $a_i\in A_{\alpha^{-1}}$, and $b_i\in A_\beta$. Then*

$$a\otimes S'_{\beta^{-1}}(b)=\sum\Delta'_{\beta,\beta^{-1}\alpha^{-1}}(a_i)(1\otimes b_i)\Leftrightarrow(1\otimes b)\Delta'_{\beta^{-1},\alpha^{-1}}(a)=\sum a_i\otimes S'^{-1}_{\beta^{-1}}(b_i).$$

2. Let $b\in A_{\alpha^{-1}},b_i\in A_\alpha,a\in A_{\alpha^{-1}\beta^{-1}}$, and $a_i\in A_{\beta^{-1}}$. Then

$$S'_{\alpha^{-1}}(b)\otimes a=\sum(b_i\otimes 1)\Delta'_{\beta^{-1}\alpha^{-1},\alpha}(a_i)\quad\Leftrightarrow\quad\Delta'_{\beta^{-1},\alpha^{-1}}(a)(b\otimes 1)=\sum S'^{-1}_{\alpha^{-1}}(b_i)\otimes a_i.$$

Lemma 3.19. *1. Let $a,a_i\in A_{\alpha^{-1}}$, $b\in A_\beta$, and $b_i\in A_{\beta^{-1}\alpha^{-1}}$. Then*

$$\Delta'_{\beta,\beta^{-1}\alpha^{-1}}(a)(1\otimes b)=\sum\Delta'_{\beta,\beta^{-1}\alpha^{-1}}(a_i)(b_i\otimes 1)\Leftrightarrow a\otimes S'^{-1}_{\beta^{-1}}(b)=\sum(1\otimes a_i)\Delta'_{\beta^{-1},\alpha^{-1}}(b_i).$$

2. Let $b \in A_{\beta^{-1}\alpha^{-1}}$ *,* $a \in A_{\alpha^{-1}}$ *,* $a_i \in A_{\alpha^{-1}}$*, and* $b_i \in A_\beta$*. Then*

$$\Delta_{\beta,\beta^{-1}\alpha^{-1}}(a)(1 \otimes b) = \sum \Delta_{\beta,\beta^{-1}\alpha^{-1}}(a_i)(b_i \otimes 1) \Leftrightarrow (1 \otimes a)\Delta_{\beta^{-1},\alpha^{-1}}(b) = \sum S_\beta(b_i) \otimes a_i.$$

The first part of the above lemma is a reformulation of lemma 3.17 for the opposite comultiplication . For part (2) if we apply σ to the equivalence in part (1), we get

$$\Delta_{\beta,\beta^{-1}\alpha^{-1}}(a)(b \otimes 1) = \sum \Delta_{\beta,\beta^{-1}\alpha^{-1}}(a_i)(1 \otimes b_i) \Leftrightarrow S_\beta(b) \otimes a = \sum (a_i \otimes 1)\Delta_{\beta^{-1},\alpha^{-1}}(b_i).$$

But this is up to a different summation gives the equivalence in part (2).

Proposition 3.20. *Let* $b \in A_\alpha, a \in A_{\alpha\beta}$*. Then*

$$(1 \otimes S_\alpha(b))\Delta_{\beta^{-1},\alpha^{-1}}(S_{\alpha\beta}(a)) = (S_\beta \otimes S_\alpha)[\Delta'_{\alpha,\beta}(a)(1 \otimes b)].$$

Proof. Write

$$\Delta_{\alpha,\beta^{-1}\alpha^{-1}}(S_\alpha(b))(1 \otimes S_{\alpha\beta}(a)) = \sum \Delta_{\beta,\beta^{-1}\alpha^{-1}}(a_i)(b_i \otimes 1).$$

By lemma 3.19 we have

$$S_\alpha(b) \otimes a = \sum (a_i \otimes 1)\Delta_{\alpha^{-1},\alpha\beta}(b_i),$$

and

$$(1 \otimes S_\alpha(b))\Delta_{\beta^{-1},\alpha^{-1}}(S_{\alpha\beta}(a)) = \sum S_\beta(b_i) \otimes a_i.$$

By lemma 3.15 we have

$$\Delta_{\alpha\beta}(a)(b \otimes 1) = \sum S_\alpha^{-1}(a_i) \otimes b_i.$$

If we combine all of this we get

$$\begin{aligned} (1 \otimes S_\alpha(b))\Delta_{\beta^{-1},\alpha^{-1}}(S_{\alpha\beta}(a)) &= \sum (S_\beta \otimes S_\alpha)(b_i \otimes S_\alpha^{-1}(a_i)) \\ &= (S_\beta \otimes S_\alpha)(\sigma(\Delta_{\alpha,\beta}(a)(b \otimes 1))) \\ &= (S_\beta \otimes S_\alpha)(\Delta'_{\alpha,\beta}(a)(1 \otimes b)). \end{aligned}$$

□

3.4 The Dual Algebra

In general, if $\Delta(A) \subset A \otimes A$ then the dual space becomes an algebra, but this is not the case in the multiplier Hopf $\pi-$coalgebra. So, in the following proposition we will use the technique of A. Van Daele in constructing the dual space.

Proposition 3.21. *If* (A, Δ) *is a multiplier Hopf* $\pi-$*coalgebra then the vector space* $A^\star$ *defined by*

$$A^\star = \oplus_{\alpha\in\pi} A^\star_\alpha,$$

where A_α^* *is the vector space of linear functionals on* A_α *spanned by elements of the form* $a \to f(bac)$ *where* $b, c \in A_\alpha$ *and* $f \in A'_\alpha$*, can be made into associative algebra under the product*

$$(fg)(a) = (f \otimes g)(\Delta_{\alpha,\beta}(a))$$

where $f \in A_\alpha^*$ *and* $g \in A_\beta^*$*.*

Proof. Let $f \in A_\alpha^*$, $g \in A_\beta^*$. Then there exist $b, c \in A_\alpha$, $d, e \in A_\beta$, $f^{'} \in A_\alpha^{'}$, and $g^{'} \in A_\beta^{'}$ such that $f(a) = f'(bac)$ for all $a \in A_\alpha$, and $g(a) = g^{'}(dae)$ for all $a \in A_\beta$. Assume that $a \in A_{\alpha,\beta}$ then

$$\begin{aligned}(fg)(a) &= m_c(f \otimes g)(\Delta_{\alpha,\beta}(a)) \\ &= (f' \otimes g')((b \otimes d)\Delta_{\alpha,\beta}(a)(c \otimes e)).\end{aligned}$$

Since $(b \otimes d)\Delta_{\alpha,\beta}(a)(c \otimes e) \in A_\alpha \otimes A_\beta$ then $(fg)(a) \in \mathbb{C}$. It means that fg is a linear functional on $A_{\alpha\beta}$. By linearity it can be extended to all $f, g \in A^*$.

By the bijectivity of $T^1_{\alpha,\beta}$ and $T^2_{\alpha,\beta}$ we can write

$$\begin{aligned}b \otimes d &= \sum_{i=1}^{n}(p_i \otimes 1)\Delta_{\alpha,\beta}(q_i), \\ c \otimes e &= \sum_{j=1}^{m}\Delta_{\alpha,\beta}(r_j)(1 \otimes k_j).\end{aligned}$$

Then

$$(fg)(a) = \sum_{i,j}(f' \otimes g')((p_i \otimes 1)\Delta_{\alpha,\beta}(q_i a r_j)(1 \otimes k_j)).$$

Define

$$h_{i,j} : A_{\alpha\beta} \to \mathbb{C}$$

by

$$h_{i,j}(a) = (f' \otimes g')((p_i \otimes 1)\Delta_{\alpha,\beta}(a)(1 \otimes k_j)).$$

Then

$$(fg)(a) = \sum_{i,j} h_{i,j}(qar).$$

i.e. $fg \in A^* \quad \forall f, g \in A^\star$ and thus $(A^*, .)$ be an algebra.

Let $f_1 \in A_\alpha^*, f_2 \in A_\beta^*$, and $f_3 \in A_\gamma^*$ and they defined by $f_i = f_i(b_i a_i c_i)$ $i = 1, 2, 3.$ where $a_1, b_1, c_1 \in A_\alpha, a_{2,}b_2, c_2 \in A_\beta$ and $a_3, b_3, c_3 \in A_\gamma$. Then

$$\begin{aligned}((f_1f_2)f_3)(a) &= (f_1f_2\otimes f_3)((1\otimes b_3)\Delta_{\alpha\beta,\gamma}(a)(1\otimes c_3))\\ &= (f_1\otimes f_2\otimes f_3)((b_1\otimes b_2\otimes 1)((1\otimes b_3)\Delta_{\alpha\beta,\gamma}(a)(1\otimes c_3))(c_1\otimes c_2\otimes 1)\\ &= (f_1\otimes f_2\otimes f_3)((1\otimes b_2\otimes b_3)((b_1\otimes 1\otimes 1)(\Delta_{\alpha,\beta}\otimes I)(\Delta_{\alpha\beta,\gamma}(a)(1\otimes c_3)))\\ &\quad (c_1\otimes c_2\otimes 1))\\ &= (f_1\otimes f_2\otimes f_3)((1\otimes b_2\otimes b_3)((I\otimes\Delta_{\beta,\alpha})((b_1\otimes 1)(\Delta_{\alpha,\beta\gamma}(a))(1\otimes 1\otimes c_3))\\ &\quad (c_1\otimes c_2\otimes 1))\\ &= (f_1\otimes f_2\otimes f_3)((1\otimes b_2\otimes b_3)((I\otimes\Delta_{\beta,\alpha})((b_1\otimes 1)(\Delta_{\alpha,\beta\gamma}(a))(c_1\otimes 1))\\ &\quad (1\otimes c_2\otimes c_3))\\ &= (f_1(f_2f_3))(a).\end{aligned}$$

Then A^* becomes an associative algebra. □

If A_α is a unital algebra $\forall\alpha\in\pi$, then we have the the same definition of the dual space introduced by Virelizier [6].

Remark 2. If $f(a)=f'(abc)$, $f\in A^*_\alpha$, and if ε is the counit of A we have

$$\begin{aligned}\varepsilon f(a) &= (\varepsilon\otimes f')((1\otimes b)\Delta_{1,\alpha}(a)(1\otimes c))\\ &= f'(b(\varepsilon\otimes I)(\Delta(a)(1\otimes c)))\\ &= f'(bac)=f(a).\end{aligned}$$

Similarly, $f\varepsilon(a)=f(a)$. So ε corresponds to the identity in the multiplier algebra of A^*.

Definition 3.22. Let A be a regular multiplier Hopf $\pi-$coalgebra and S is the antipode of A. Then we define the adjoint antipode to be

$$S^\star : A^*\to A^*$$

by

$$(S^\star f)(a)=f(S_{\alpha^{-1}}(a))$$

for all $f\in A^*_\alpha, a\in A_{\alpha^{-1}}$.

Lemma 3.23. *$S^\star f\in A^*$ if $f\in A^*$.*

Proof. Let $f(a)=f^{'}(bac)$ where $f\in A^*_\alpha, a,b,c\in A_{\alpha^{-1}}$, and choose $d\in A_1$ such that $\varepsilon(d)=1$. By using theorem 2 we get

$$\begin{aligned}(S^{\star}f)(a) &= f(S_{\alpha^{-1}}(a))=f'(bS_{\alpha^{-1}}(a)c)\\ &= f'(b\varepsilon_\alpha(d)S_{\alpha^{-1}}(a)c)\\ &= f'(m_\alpha((I\otimes S_{\alpha^{-1}})((b\otimes 1)\Delta_{\alpha,\alpha^{-1}}(d))(1\otimes S_{\alpha^{-1}}(a)c))).\end{aligned}$$

By the bijectivity of $T^2_{\alpha,\alpha^{-1}} \exists\ p_i \otimes q_i \in A_\alpha \otimes A_{\alpha^{-1}}$ such that $(b \otimes 1)\Delta_{\alpha,\alpha^{-1}}(d) = \sum_{i=1}^n p_i \otimes q_i$. Then

$$\begin{aligned}(S^\star f)(a) &= \sum f'(m_\alpha(p_i \otimes S_{\alpha^{-1}}(aq_i)c) \\ &= \sum f'(p_i S_{\alpha^{-1}}(aq_i)c).\end{aligned}$$

By a similar argument we can prove that

$$(S^\star f)(a) = \sum f^{'}(b_i S_{\alpha^{-1}}(r_j a q_i)k_j).$$

□

Lemma 3.24. *If (A, Δ) is a regular multiplier Hopf π- coalgebra then $S^\star$ is an antihomomorphism of A^*.*

Proof. Let $f \in A^*_{\beta^{-1}}, g \in A^*_{\alpha^{-1}}$, and write $g(a_1) = g'(S_\alpha(b)a_1)$ for some $b \in A_\alpha$. Then for $a \in A_{\alpha\beta}$ we have

$$\begin{aligned}S^\star(fg)(a) &= fg(S_{\alpha\beta}(a)) = (f \otimes g')((1 \otimes S_\alpha(b))\Delta_{\beta^{-1},\alpha^{-1}}(S_{\alpha\beta}(a))) \\ &= (f \otimes g')((S_\beta \otimes S_\alpha)(\Delta'_{\alpha\beta}(a)(1 \otimes b))) \\ &= (S^\star g' \otimes S^\star f)(\Delta(a)(b \otimes 1)).\end{aligned}$$

But

$$S^\star g'(ab) = g'(S_\alpha(b)S_\alpha(a)) = g(S_\alpha(a)) = S^\star g(a),$$

then

$$S^\star(fg) = ((S^\star g)(S^\star f))(a).$$

□

Lemma 3.25. *Let $A = \{A_p\}_{p\in G}$ be a multiplier Hopf $G-$coalgebra. Then $H = \{p \in G : A_p \neq 0\}$ is a subgroup of G.*

Proof. Let $H = \{p \in G : A_p \neq 0\}$ and $p \in H$. Then from the bijectivity of the map $T^1_{1,p} : A_p \otimes A_p \to A_1 \otimes A_p$ we get that $A_1 \neq 0$. Also, if $p, q \in H$ then the bijectivity of the map $T^1_{p,q} : A_{p,q} \otimes A_q \to A_p \otimes A_q$ implies that $A_{p,q} \neq 0$. Finally, for all $p \in H$ the bijectivity of the map $T^1_{p^{-1},p} : A_1 \otimes A_p \to A_{p^{-1}} \otimes A_p$ implies that $p^{-1} \in H$. □

4 The second approach

Now let $(A = \{A_p\}_{p\in G}, \Delta = \{\Delta_{p,q}\}_{p,q\in G})$ be a multiplier Hopf $G-$coalgebra . Let A_G be the direct sum of the algebras A_p. An element in A_G is a function a from G to $\cup A_p$, with finite support and so that $a(p) \in A_p\ \forall p \in G$. This algebra is an algebra with a nondegenerate product. For $a \in A_p$ we will use a for the function in A_G that have the value a at p and 0 otherwise and use $a(p)$ when we mean the element $a \in A_p$.

Proposition 4.1. *let* $(A = \{A_p\}_{p\in G}, \Delta = \{\Delta_{pq}\}_{p,q\in G})$ *be a multiplier Hopf* $G-$*coalgebra . Then* A_G *becomes a multiplier Hopf algebra under the comultiplication* $\Delta_G : A_G \to M(A_G \otimes A_G)$ *given by*

$$\Delta_G(a)(p, q) = \Delta_{p,q}(a(pq)).$$

Moreover there exist a nondegenerate homomorphism $\gamma : K(G) \to M(A_G)$ *,where* $K(G)$ *be the algebra of complex valued functions on* G *with finite support, such that* $\gamma(K(G))$ *is in the center of* $M(A_G)$*. And* γ *is compatible with the comultiplication.*

Proof. $I-$ (A_G, Δ_G) is a multiplier Hopf algebra.
(1) Let $a, b \in A_G$. Then

$$(\Delta_G(a)(1 \otimes b))(p, q) = \Delta_{p,q}(a(pq))(1 \otimes b(q)),$$

but b has a finite support, then $\Delta_G(a)(1 \otimes b) \in A_G \otimes A_G$. Similarly, $(a \otimes 1)\Delta_G(b) \in A_G \otimes A_G$.
(2) Let $a, b, c \in A_G$. Then we have

$$\begin{aligned}(a \otimes 1 \otimes 1)(\Delta_G \otimes I)(\Delta_G(b)(1 \otimes c))(p, q, r) &= (a(p) \otimes 1 \otimes 1)(\Delta_{p,q} \otimes I)[(\Delta_G(b)(1 \otimes c))(pq, r)]\\ &= (a(p) \otimes 1 \otimes 1)(\Delta_{p,q} \otimes I)[\Delta_{pq,r}(b(pqr))(1 \otimes c(r))]\\ &= (I \otimes \Delta_{q,r})[(a(p) \otimes 1)\Delta_{p,qr}(b(pqr))](1 \otimes 1 \otimes c(r))\\ &= [(I \otimes \Delta_G)[(a \otimes 1)\Delta_G(b))](1 \otimes 1 \otimes c)](p, q, r).\end{aligned}$$

(3) The map $T_1 : A \otimes A \longrightarrow A \otimes A$ defined by

$$T_1(a \otimes b)_{(p,q)} = \Delta_G(a)(1 \otimes b)(pq) = \Delta_{p,q}(a(pq))(1 \otimes b(q)) = T^1_{p,q}(a(pq) \otimes b(q))$$

is a bijective with the inverse

$$T_1^{-1}(a \otimes b)(p, q) = (T^1_{p,q})^{-1}(a(pq^{-1}) \otimes b(q)),$$

since

$$\begin{aligned}((T_1^{-1} \circ T_1)(a \otimes b))(p, q) &= (T^1_{p,q})^{-1}(T_1(a \otimes b)(pq^{-1}, q))\\ &= (T^1_{p,q})^{-1}(T^1_{p,q}(a(pq^{-1}q) \otimes b(q))\\ &= a(p) \otimes b(q).\end{aligned}$$

Similarly, $[(T_1 \circ T_1^{-1}(a \otimes b)](p, q) = a(p) \otimes b(q)$.
Also, the map $T_2 : A \otimes A \longrightarrow A \otimes A$ defined by

$$T_2(a \otimes b)(p, q) = (a(p) \otimes 1)\Delta_{p,q}(b(pq))$$

is bijective with the inverse

$$T_2^{-1}(a \otimes b)(p, q) = (T^2_{p,q})^{-1}(a(p) \otimes b(p^{-1}q)).$$

$II-$ There exist a nondegenerate homomorphism γ, from $K(G)$ into the center of $M(A_G)$, compatible with the comultiplication.
(1) Let $K(G)$ be the algebra of complex valued functions on the group G with finite support. Define

$$\gamma : K(G) \longrightarrow M(A)$$

by

$$\gamma(f)(p) = f(p)1_p$$

By the definition of γ we have that $\gamma(f)$ in the center $M(A)$ $\forall f \in K(G)$. Moreover, $\gamma(fg)(p) = (fg)(p)1_p = f(p)g(p)1_P = (\gamma(f)\gamma(g))(p)$ and $\gamma(K(G))A_G = A_G\gamma(K(G)) = A_G$.
(2) Consider

$$\begin{aligned}(\gamma \otimes \gamma)\Delta_\circ(f)(p,q) &= [(\gamma \otimes \gamma)(\sum_s C_s \sum_r \delta_r \otimes \delta_{r^{-1}s}(p,q))] \\ &= [\sum_{s,r\in G} C_s\delta_r(p) \otimes \delta_{r^{-1}s}(q)1_q] \\ &= f(pq)(1_p \otimes 1_q).\end{aligned}$$

On the other hand

$$\begin{aligned}\Delta_G(\gamma(f))(p,q) &= \Delta_{p,q}(\gamma(f)(p,q)) \\ &= \Delta_{p,q}(f(pq)1_{p,q}) \\ &= f(pq)(1_p \otimes 1_q).\end{aligned}$$

Thus γ is compatible with the comultiplication □

Let (A, Δ) be a multiplier Hopf algebra such that there exist a nondegenerate homomorphism $\gamma : K(G) \longrightarrow M(A)$ such that $\gamma(K(G))$ is in the center of $M(A)$ and

$$\Delta(\gamma(f)) = (\gamma \otimes \gamma)\Delta_\circ(f),$$

where $\Delta_\circ(f)(p,q) = f(pq)$.
We denote this multiplier Hopf algbera with the nondegenerate homomorphism γ by $(\tilde{A}, \tilde{\Delta})$.

Let $A_p = \tilde{A}\gamma(\delta_p)$, $\delta_p \in K(G)$ where $\delta_p(q) = \delta_p^q = \{^{0,\ p\neq q}_{1,\ p=q}$. Because of A_p is a subalgebra of $\tilde{A}$ and satisfies that if $a\gamma(\delta_p)b\gamma(\delta_p) = 0\ \forall\ a\gamma(\delta_p), \gamma(\delta_p) \neq 0$ then $ab\gamma(\delta_p) = 0\ \forall\ a \in \tilde{A}$ and hence $b\gamma(\delta_p) = 0$. Similarly, if $a\gamma(\delta_p)b\gamma(\delta_p) = 0\ \forall\ b\gamma(\delta_p), \gamma(\delta_p) \neq 0$ then $a\gamma(\delta_p) = 0$. Then A_p is an algebra with a nondegenerate product. $\tilde{A}$ can be identified with the direct of A_p's using

$$a \longrightarrow (a\gamma(\delta_p))_{p\in G}$$

Since γ is a nondegenerate homomorphism then for all $a \in \tilde{A}$ there exist $f \in K(G)$ such that $a = a\gamma(f)$. Since $f \in K(G)$ then there exist finite number of $\delta's$ such that $f = \sum c_p\delta_p$ and hence $a \in \oplus_p A\gamma(\delta_p) = \oplus_p A_p$. Then $\forall a \in \tilde{A}$ there exist a finite numbers of p's such that $a \in \oplus_p A_p$

Let $a, b \in \tilde{A}$ and $\delta_p, \delta_q \in K(G)$. Then the map $\tilde{T}_1$ satisfies

$$\begin{aligned}
\tilde{T}_1(a\gamma(\delta_p) \otimes b\gamma(\delta_q)) &= \tilde{\Delta}(a\gamma(\delta_p))(1 \otimes b\gamma(\delta_q)) \\
&= \tilde{\Delta}(a)(\gamma \otimes \gamma)\Delta_\circ(\delta_p)(1 \otimes b)(1 \otimes \gamma(\delta_q)) \\
&= \tilde{\Delta}(a)(1 \otimes b)(\gamma \otimes \gamma)T_1^\circ(\delta_p \otimes \delta_q) \\
&= \tilde{T}_1(a \otimes b)(\gamma \otimes \gamma)T_1^\circ(\delta_p \otimes \delta_q)
\end{aligned}$$

and its inverse satisfies

$$\tilde{T}_1^{-1}(a\gamma(\delta_p) \otimes b\gamma(\delta_q)) = \tilde{T}_1^{-1}(a \otimes b)(\gamma \otimes \gamma)(T_1^\circ)^{-1}(\delta_p \otimes \delta_q).$$

Similarly, the map $\tilde{T}_2$ satisfies

$$\tilde{T}_2(a\gamma(\delta_p) \otimes b\gamma(\delta_q)) = \tilde{T}_2(a \otimes b)(\gamma \otimes \gamma)T_2^\circ(\delta_p \otimes \delta_q)$$

and its inverse satisfies

$$\tilde{T}_2^{-1}(a\gamma(\delta_p) \otimes b\gamma(\delta_q)) = \tilde{T}_2^{-1}(a \otimes b)(\gamma \otimes \gamma)(T_2^\circ)^{-1}(\delta_p \otimes \delta_q).$$

For $a, b \in \tilde{A}$ and $\delta_p, \delta_q \in K(G)$ the counit satisfies

$$\begin{aligned}
\tilde{\varepsilon}(a\gamma(\delta_p)b\gamma(\delta_q)) &= m\tilde{T}_1^{-1}(a\gamma(\delta_p) \otimes b\gamma(\delta_q)) \\
&= m\tilde{T}_1^{-1}(a \otimes b)(\gamma \otimes \gamma)((T_1^\circ)^{-1}(\delta_p \otimes \delta_q)) \\
&= \tilde{\varepsilon}(a)b\gamma(\varepsilon_\circ(\delta_p)\delta_q) \\
&= \tilde{\varepsilon}(a)\varepsilon_\circ(\delta_p)b\gamma(\delta_q).
\end{aligned}$$

Since $\varepsilon_\circ(f) = f(e) \;\; \forall f \in K(G)$ then $\tilde{\varepsilon}(a\gamma(\delta_p)) = \tilde{\varepsilon}(a)$ when $e = p$ and zero otherwise
Also, the counit satisfies

$$(I \otimes \tilde{\varepsilon})\tilde{\Delta}(a\gamma(\delta_p))(1 \otimes b\gamma(\delta_q)) = a\gamma(\delta_p)b\gamma(\delta_q) = ab\gamma(\delta_p\delta_q),$$

$$(\tilde{\varepsilon} \otimes I)(a\gamma\delta_p \otimes 1)\tilde{\Delta}(b\gamma\delta_q) = ab\gamma(\delta_p)\gamma(\delta_q) = ab\gamma(\delta_p\delta_q).$$

For the antipode we have

$$\begin{aligned}
\tilde{S}(a\gamma(\delta_p))b\gamma(\delta_p) &= (\tilde{\varepsilon} \otimes I)\tilde{T}_1^{-1}(a\gamma(\delta_p) \otimes b\gamma(\delta_q)) \\
&= (\tilde{\varepsilon} \otimes I)\tilde{T}_1^{-1}(a \otimes b)(\gamma \otimes \gamma)((T_1^\circ)^{-1}(\delta_p \otimes \delta_q)) \\
&= \tilde{S}(a)b\gamma((\varepsilon_\circ \otimes I)(T_1^\circ)^{-1}(\delta_p \otimes \delta_q)) \\
&= \tilde{S}(a)b\gamma(S_\circ(\delta_p)\gamma(\delta_q) \\
&= \tilde{S}(a)\gamma(S_\circ(\delta_p))b\gamma(\delta_q).
\end{aligned}$$

Since $S_\circ(\delta_p) = \delta_{p^{-1}}$ then $\tilde{S}(a\gamma(\delta_p)) = \tilde{S}(a)\gamma(\delta_{p^{-1}})$.
Also, the antipode satisfies

$$\begin{aligned} m(S \otimes I)\Delta(a\gamma(\delta_p))(1 \otimes b\gamma(\delta_q)) &= \varepsilon(a\gamma(\delta_p))b\gamma(\delta_q) = \varepsilon(a)\varepsilon_\circ(\delta_p)b\gamma(\delta_q), \\ m(I \otimes S)(b\gamma(\delta_q) \otimes 1)\Delta(a\gamma(\delta_p) &= b\gamma(\delta_q)\varepsilon(a)\varepsilon_\circ(\delta_p). \end{aligned}$$

Proposition 4.2. *Let $\tilde{A}$ be a multiplier Hopf algebra and $\gamma : K(G) \longrightarrow M(\tilde{A})$ be a nondegenerate homomorphism from $K(G)$ into the center of $M(\tilde{A})$ satisfies*

$$\tilde{\Delta}(\gamma(f)) = (\gamma \otimes \gamma)\Delta_\circ(f)$$

then this comes from a multiplier Hopf $G-$coalgebra.

Proof. Define a family of algebras $A = \{A_P\}_{p\in G}$ by $A_p = \tilde{A}\gamma(\delta_p)$. Then A_p is an algebra with nondegenerate product. Define a family of homomorphism $\Delta = \{\Delta_{p,q} : A_{pq} \longrightarrow M(A_p \otimes A_q)\}_{p,q\in G}$ by

$$\Delta_{p,q}(a\gamma(\delta_{pq})) = \tilde{\Delta}(a)(\gamma(\delta_p) \otimes \gamma(\delta_q)).$$

For $a\gamma(\delta_{pq}) \in A_{pq}, b\gamma(\delta_q) \in A_q$ we have

$$\begin{aligned} \Delta_{p,q}(a\gamma(\delta_{pq}))(1 \otimes b\gamma(\delta_q)) &= \tilde{\Delta}(a)(1 \otimes b)(\gamma(\delta_p) \otimes \gamma(\delta_q)) \\ &= \tilde{T}_1(a \otimes b)(\gamma(\delta_p) \otimes \gamma(\delta_q)) \in A_p \otimes A_q. \end{aligned}$$

For $b\gamma(\delta_{pq}) \in A_{pq}$ and $a\gamma(\delta_p) \in A_p$ we have

$$\begin{aligned} (a\gamma(\delta_p) \otimes 1)\Delta_{pq}(b\gamma(\delta_{pq})) &= (a \otimes 1)\tilde{\Delta}(b)(\gamma(\delta_p) \otimes \gamma(\delta_q)) \\ &= \tilde{T}_2(a \otimes b)(\gamma(\delta_p) \otimes \gamma(\delta_q)) \in A_p \otimes A_q. \end{aligned}$$

The coassociativity of $\Delta = \{\Delta_{p,q} : A_{pq} \longrightarrow M(A_p \otimes A_q)\}_{p,q\in G}$ is follows directly because of the coassociativity of $\tilde{\Delta}$ and that $\gamma(K(G))$ is in the center of $M(A)$. So Δ is a comultiplication
The maps $T^1_{p,q} : A_{pq} \otimes A_q \longrightarrow A_p \otimes A_q$ given by

$$\begin{aligned} T^1_{p,q}(a\gamma(\delta_p) \otimes b\gamma(\delta_q)) &= \Delta_{p,q}(a\gamma(\delta_{pq})(1 \otimes b\gamma(\delta_q))) \\ &= \tilde{T}_1(a \otimes b)(\gamma(\delta_p) \otimes \gamma(\delta_q)) \\ &= \tilde{T}_1(a \otimes b)(\gamma \otimes \gamma)(T^1_\circ(\delta_{pq} \otimes \delta_q)) \end{aligned}$$

are bijective with inverses

$$\begin{aligned} (T^1_{p,q})^{-1}(a\gamma(\delta_p) \otimes b\gamma(\delta_q)) &= \tilde{T}_1^{-1}(a \otimes b)(\gamma(\delta_{pq}) \otimes \gamma(\delta_q)) \\ &= \tilde{T}_1^{-1}(a \otimes b)(\gamma \otimes \gamma)T^1_\circ(\delta_p \otimes \delta_q) \end{aligned}$$

Also, the maps $T^2_{p,q} : A_p \otimes A_{pq} \longrightarrow A_p \otimes A_q$ given by

$$\begin{array}{rcl} T^2_{p,q}(a\gamma(\delta_p) \otimes b\gamma(\delta_q)) & = & (a\gamma(\delta_p) \otimes 1)\Delta_{p,q}(a\gamma(\delta_{pq})) \\ & = & \tilde{T}_2(a \otimes b)(\gamma \otimes \gamma)T^2_\circ(\delta_p \otimes \delta_{pq}) \end{array}$$

are bijective with inverses

$$\begin{array}{rcl} (T^2_{p,q})^{-1}(a\gamma(\delta_p) \otimes b\gamma(\delta_q)) & = & \tilde{T}_2^{-1}(a \otimes b)(\gamma(\delta_p) \otimes \gamma(\delta_{pq})) \\ & = & \tilde{T}_2^{-1}(a \otimes b)(\gamma \otimes \gamma)((T^2_\circ)^{-1}(\delta_p \otimes \delta_{pq})) \end{array}$$

Thus $(A = \{A_p\}_{p\in G}, \Delta = \{\Delta_{p,q}\}_{p,q\in G})$ is a multiplier Hopf $G-$coalgebra.

Now let (A_G, Δ_G) be the multiplier Hopf algebra associated to (A, Δ) as in lemma(1.4). Define a map $\varphi : \tilde{A} \longrightarrow A_G$ by

$$\varphi(a)(p) = a\gamma(\delta_p) \in A_p.$$

Since $a \in \oplus_p A_p$ for finite number of p's then $\varphi(a)$ will has finite support (i.e. φ is well defined $(\varphi(\tilde{A}) \subset A_G)$).
If $\varphi(a) = \varphi(b),\ \ a, b \in \tilde{A}$ then

$$a\gamma(\delta_p) = b\gamma(\delta_p)\ \forall p \Longrightarrow (a - b)\gamma(\delta_p) = 0\ \forall \delta_p$$

By the nondegenerancy of γ we have $a - b = 0 \Longrightarrow a = b$ hence φ is $1 - 1$.
Also,

$$\forall\ b \in A_G\ \exists\ a = \sum_{p\in G} b(p) \in \tilde{A} \text{ such that } \varphi(a) = b,$$

then φ is onto.
φ is a nondegenerate homomorphism, where

$$\varphi(ab)(p) = ab\gamma(\delta_p) = a\gamma(\delta_p)b\gamma(\delta_p) = (\varphi(a)\varphi(b))(p)\ \forall\ p \in G.$$

Moreover, we also have

$$\begin{array}{rcl} \Delta_G(\varphi(a))(p, q) & = & \Delta_{p,q}(\varphi(a)(pq)) = \Delta_{p,q}(a\gamma(\delta_{pq})) \\ & = & \tilde{\Delta}(a)(\gamma(\delta_p) \otimes \gamma(\delta_q)) = ((\varphi \otimes \varphi)\Delta(a))(p, q) \end{array}$$

Thus $\tilde{A}$ is isomorphic to A_G.
There exist a map $\gamma_G : K(G) \longrightarrow A_G$ defined by

$$\gamma_G(f)(p) = f(p)\gamma(\delta_p).$$

Since $\gamma(\delta_p)$ is the unit of $M(A_p)$. Then $\gamma_G(f)(p) = f(p)1_p$ and hence as in lemma (1.4) γ_G is a nondegenerate homomorphism compatible with the comultiplication and $\gamma_G(K(G))$ is in the central of $M(A)$.

□

The last two propositions show that there exist a one to one correspondence between the class of multiplier Hopf $G-$coalgebras and the class of multiplier Hopf algebras such that each multiplier Hopf algbera endowed with a nondegenerate hommomophism form the algebra of complex valued functions with finite support on the group G into the center of its multiplier algebra and compatible with its comultiplication.

Lemma 4.3. *Let (A, Δ) be a multiplier Hopf $G-$coalgebra. Then there exist a homomorphism $\varepsilon : A_e \longrightarrow \mathbb{C}$ such that*

$$(I \otimes \varepsilon)((a \otimes 1)\Delta_{p,e}(b)) = ab,$$

$$(\varepsilon \otimes I)(\Delta_{e,p}(a)(1 \otimes b)) = ab$$

for all $a, b \in A_p,\ p \in G$.

Proof. Consider the algebra $(\tilde{A}, \tilde{\Delta})$ be as above. For $a, b \in \tilde{A}$ we have

$$\varepsilon(a)b = \sum_p \sum_q \varepsilon(a\gamma(\delta_p))b\gamma(\delta_q) = \varepsilon(a\gamma(\delta_e))b.$$

Define a map $\varepsilon : A_e \to \mathbb{C}$ by $\varepsilon(a) = \tilde{\varepsilon}(a)\ \ \forall a \in A_e$. By the properties of the map $\tilde{\varepsilon}$ we have that ε is a nondegenerate homomorphism. Let $x, y \in A_p$ with $x = a\gamma(\delta_p), y = b\gamma(\delta_p)$ where $a, b \in \tilde{A}$ then

$$\begin{aligned}
(I \otimes \varepsilon)((x \otimes 1)\Delta_{p,e}(y)) &= (I \otimes \varepsilon)((a\gamma(\delta_p) \otimes 1)\Delta_{p,e}(b\gamma(\delta_p))) \\
&= (I \otimes \tilde{\varepsilon})((a \otimes 1)(\gamma \otimes \gamma)(\delta_p \otimes \delta_e)\tilde{\Delta}(b)) \\
&= (I \otimes \tilde{\varepsilon})((a \otimes 1)(\gamma \otimes \gamma)((\delta_p \otimes 1)\Delta_\circ(\delta_p))\tilde{\Delta}(b)) \\
&= (I \otimes \tilde{\varepsilon})((a\gamma(\delta_p) \otimes 1)\tilde{\Delta}(b\gamma(\delta_p))) \\
&= a\gamma(\delta_p)b\gamma(\delta_p) = xy.
\end{aligned}$$

Similarly, we can prove that $(\varepsilon \otimes I)(\Delta_{e,p}(x)(1 \otimes y)) = xy$.
Also, for the multiplier Hopf algebra (A_G, Δ_G) we get that $\varepsilon_G(a) = \varepsilon(a(e))\ \ \forall a \in A_G$. ☐

Lemma 4.4. *If (A, Δ) is a multiplier Hopf $G-$coalgebra then there exist a family of antihomomorphisms $S = \{S_p : A_p \longrightarrow M(A_{p^{-1}})\}_{p \in G}$ such that for all $a \in A_e,\quad b \in A_p$ and $\phi \in G$ we have*

$$m_p((I \otimes S_{p^{-1}})((b \otimes 1)\Delta_{p,p^{-1}}(a))) = \varepsilon(b)a,$$

$$m_p((S_{p^{-1}} \otimes I)\Delta_{p^{-1},p}(a)(1 \otimes b)) = \varepsilon(a)b.$$

Proof. In the algebra $(\tilde{A}, \tilde{\Delta})$

$$\tilde{S}(a\gamma(\delta_p)) = \tilde{S}(a)\gamma(S_\circ(\delta_p)).$$

For the multiplier Hopf $G-$coalgebra (A, Δ) we define

$$S_p : A_p \longrightarrow M(A_{p^{-1}})$$

by

$$S(a\gamma(\delta_p)) = S(a)\gamma(S_\circ(\delta_p)).$$

Let $a \in A_e$ $b \in A_p$. Then

$$\begin{aligned} m_p(I \otimes S_{p^{-1}})[(b\gamma(\delta_p) \otimes 1)\Delta_{p,p^{-1}}(a\gamma(\delta_e))] &= m(I \otimes S)[(b \otimes 1)\tilde{\Delta}(a)(\gamma \otimes \gamma)(\delta_p \otimes 1)\Delta_\circ(\delta_e)] \\ &= m(I \otimes S)[(b\gamma(\delta_p) \otimes 1)\tilde{\Delta}(a\gamma(\delta_e))] \\ &= \varepsilon(a\gamma(\delta_e))b\gamma(\delta_p). \end{aligned}$$

Also, we can prove that

$$m_p(S_{p^{-1}} \otimes I)(\Delta_{p^{-1},p}(a)(1 \otimes b)) = \varepsilon(a)b.$$

For the algebra (A_G, Δ_G), we have

$$S(a)(p) = S_{p^{-1}}(a(p^{-1})).$$

□

Theorem 4. *A is a Hopf $G-$coalgebra if and only if A is multiplier Hopf $G-$coalgebra such that A_p is unital $\forall p \in G$.*

Proof. If (A, Δ) is a Hopf $G-$coalgebra then A_p has an identity $\forall p \in G$ which means that $M(A_p) = A_p \quad \forall p \in G$ and according to remark (1) we have Δ is a comultiplication.
Now we will prove that the linear maps

$$\begin{aligned} T^1_{p,q} &: A_{pq} \otimes A_q \longrightarrow A_p \otimes A_q \quad , \\ T^2_{p,q} &: A_p \otimes A_{pq} \longrightarrow A_p \otimes A_q \end{aligned}$$

defined by

$$\begin{aligned} T^1_{p,q}(a \otimes b) &= \Delta_{p,q}(a)(1 \otimes b) \quad , \\ T^2_{p,q}(a \otimes b) &= (a \otimes 1)\Delta_{p,q}(b) \end{aligned}$$

are bijective for all $p, q \in G$.
Consider the following linear maps

$$\begin{aligned} R^1_{p,q} &: A_p \otimes A_q \longrightarrow A_{pq} \otimes A_q \quad , \\ R^2_{p,q} &: A_p \otimes A_q \longrightarrow A_p \otimes A_{pq} \end{aligned}$$

which defined by

$$\begin{aligned} R^1_{p,q}(a\otimes b) &= ((I\otimes S_{q^{-1}})\Delta_{pq,q^{-1}}(a))(1\otimes b) \quad , \\ R^2_{p,q}(a\otimes b) &= (a\otimes 1)((S_{p^{-1}}\otimes I)\Delta_{p^{-1},pq}(b)) \quad . \end{aligned}$$

By using the properties of S and Δ we can prove that $R^1_{p,q}$ and $R^2_{\phi,q}$ are the inverses of $T^1_{\phi,q}$ and $T^2_{\phi,q}$, respectively. By using Sweedler's notation, we get

$$\begin{aligned} T^1_{\phi,q}R^1_{\phi,q}(a\otimes b) &= \sum_{(a)} T^1_{\phi,q}(a_{(1,\phi q)}\otimes S_{q^{-1}}(a_{(2,q^{-1})})\, b) \\ &= \sum_{(a)} a_{(1,\phi)}\otimes a_{(2,q)}S_{q^{-1}}(a_{(3,q^{-1})})\, b \\ &= \sum_{(a)} a_{(1,\phi)}\otimes m_\phi(I\otimes S_{q^{-1}})\Delta_{q,q^{-1}}(a_{(2,1)})(1\otimes b) \\ &= \sum_{(a)} (a_{(1,\phi)}\otimes \varepsilon(a_{(2,1)})1_q)(1\otimes b) = \, a\otimes b. \end{aligned}$$

Similarly, for $R^1_{\phi,q}T^1_{\phi,q}$, $R^2_{\phi,q}T^2_{\phi,q}$ and $T^2_{\phi,q}R^2_{\phi,q}$. Therefore A is a multiplier Hopf $\pi-$coalgebra.

If A is a multiplier Hopf $G-$coalgebra such that A_p is unital $\forall p\in G$ then $M(A_\phi\otimes A_q)=A_\phi\otimes A_q\ \forall\, \phi,q\in G$ and by using remark (1) we have

$$(\Delta_{\phi,q}\otimes I)\Delta_{\phi q,r} = (I\otimes\Delta_{q,r})\Delta_{\phi,qr}$$

as a maps from $A_{\phi qr}$ into $A_\phi\otimes A_q\otimes A_r$.
Since A is unital then $M(A_1\otimes A_\phi)=A_1\otimes A_\phi$ and $M(A_\phi\otimes A_1)=A_\phi\otimes A_1$ so the counitary property just means

$$(I\otimes\varepsilon)\Delta_{\phi,1} = (\varepsilon\otimes 1)\Delta_{1,\phi} = I \quad \forall\phi\in G.$$

Then we obtain the definition of the counit introduced by Turaev.
If A is unital then the antihomomorphism S has the form

$$S=\{S_\phi : A_\phi \longrightarrow A_{\phi^{-1}}\}_{\phi\in G}$$

such that

$$m_\phi(S_{\phi^{-1}}\otimes I)\Delta_{\phi^{-1},\phi} = \varepsilon 1_\phi = m_\phi(I\otimes S_{\phi^{-1}})\Delta_{\phi,\phi^{-1}}.$$

□

4.1 Regular multiplier Hopf group coalgebra.

Lemma 4.5. *$(A=\{A_p\}_{p\in G},\Delta)$ is a regular multiplier Hopf $G-$coalgebra iff (A_G,Δ_G) is a regular multiplier Hopf algbera .*

Proof. 1) If $(A = \{A_p\}_{p\in G}, \Delta)$ is a regular multiplier Hopf $G-$coalgebra , then (A_G, Δ_G) is a multiplier Hopf algbera . It is clear that Δ_G is nondegenerate and from the coassociativity of Δ_G, it follows that Δ'_G is coassociative. Let $a, b \in A_G$ then

$$\begin{aligned}\Delta'_G(a)(1\otimes b)(q,p) &= \sigma(\Delta_G(a)(b\otimes 1))(q,p)\\ &= \sigma(\Delta_G(a)(b\otimes 1)(p,q))\\ &= \sigma(\Delta_{p,q}(a(pq)(b(p)\otimes 1)))\\ &= \Delta'_{p,q}(a(pq)(1\otimes b(p))) \in A_q\otimes A_p.\end{aligned}$$

Similarly, we can prove that $(a\otimes 1)\Delta'_G(b) \in A_q \otimes A_p$. Hence, Δ_G is an opposite comultiplication on A_G.

The map $T'_1 : A_G \otimes A_G \rightarrow A_G \otimes A_G$ defined by

$$\begin{aligned}T'_1(a\otimes b)(q,p) &= \Delta'_G(a)(1\otimes b)(q,p)\\ &= \Delta'_{p,q}(a(pq)(1\otimes b(p)))\\ &= T'^{1}_{p,q}(a(pq)(1\otimes b(p))) \in A_q\otimes A_p\end{aligned}$$

is bijective since the members of the family $T'^{1} = \{T'^{1}_{p,q}\}_{p,q\in G}$ are bijective .

Similarly, we can prove that the map T'_2 is bijective.

Therefor, (A_G, Δ_G) is a regular multiplier Hopf algbera .

2)If (A_G, Δ_G) is a regular multiplier Hopf algbera , then (A, Δ) is a multiplier Hopf $G-$coalgebra .

a)Let $a \in A_{pq}, b \in A_p$. Then

$$\begin{aligned}\Delta'_G(a)(1\otimes b)(q,p) &= \sigma_G(\Delta(a)(b\otimes 1)(p,q))\\ &= \sigma(\Delta(a)(b\otimes 1)(p,q))\\ &= \sigma(\Delta(a(pq))(b(p)\otimes 1))\\ &= \Delta'_{p,q}(a(pq))(1\otimes b(p)).\end{aligned}$$

Since $\Delta'_G(a)(1\otimes b)(q,p) \in A_q\otimes A_p$ then $\Delta'_{p,q}(a)(1\otimes b) \in A_q\otimes A_p$.

Similarly, we can prove that $(a\otimes 1)\Delta'_{p,q}(b) \in A_q\otimes A_p$ for all $a\in A_q, b\in A_{pq}$

b) Since the map T'_1 is bijective then $T'_1(p,q)$ is bijective . Similarly, for the maps $T'^{2}_{p,q}$.

c) Let $a \in A_p, b\in A_{rqp}$ and $c \in A_r$. Then

$$\begin{aligned}[(a\otimes 1\otimes 1) \quad & (\Delta'_G\otimes I)(\Delta'_G(b)(1\otimes c))](p,q,r)\\ = \quad & [(I\otimes\sigma_G)(\sigma_G\otimes I)(I\otimes\sigma_G)((1\otimes 1\otimes a)(I\otimes\Delta_G)(\Delta_G(b)(c\otimes 1)))](p,q,r)\\ = \quad & (I\otimes\sigma)(\sigma\otimes I)(I\otimes\sigma)((1\otimes 1\otimes a)(I\otimes\Delta_G)(\Delta_G(b)(c\otimes 1))(r,q,p))\\ = \quad & (I\otimes\sigma)(\sigma\otimes I)(I\otimes\sigma)(1\otimes 1\otimes a(p))(I\otimes\Delta_{q,p})(\Delta_{r,qp}(b(rqp)(c(r)\otimes 1)))\\ = \quad & (a(p)\otimes 1\otimes 1)(\Delta'_{q,p}\otimes I)(\Delta'_{r,qp}(b(rqp))(1\otimes c(r))).\end{aligned}$$

On the other hand

$$\begin{aligned}(I \otimes \Delta'_G) \quad & ((a \otimes 1)\Delta'_G(b))(1 \otimes 1 \otimes c)(p,q,r) \\ = \quad & (\sigma_G \otimes I)(I \otimes \sigma_G)(\sigma_G \otimes I)((\Delta_G \otimes I)((1 \otimes a)\Delta_G(b))(c \otimes 1 \otimes 1))(p,q,r) \\ = \quad & \sigma \otimes I)(I \otimes \sigma)(\sigma \otimes I)((\Delta_G \otimes I)((1 \otimes a)\Delta_G(b))(c \otimes 1 \otimes 1)(r,q,p)) \\ = \quad & (\sigma \otimes I)(I \otimes \sigma)(\sigma \otimes I)((\Delta_{r,q} \otimes I)((1 \otimes a(p))\Delta_{rq,p}(b(rq,p)))(c(r) \otimes 1 \otimes 1)) \\ = \quad & (I \otimes \Delta'_{r,q})((a(p) \otimes 1)\Delta'_{rq,p}(b(rq,p)))(1 \otimes 1 \otimes c(r)).\end{aligned}$$

Since Δ'_G is coassociative, we have

$$(a(p)\otimes 1\otimes 1)(\Delta'_{q,p}\otimes I)(\Delta_{r,qp}(b(r,qp))(1\otimes c(r))) = (I\otimes\Delta'_{r,q})((a(p)\otimes 1)\Delta_{rq,p}(b(rq,p)))(1\otimes 1\otimes c(r)).$$

Then (A,Δ) is regular multiplier Hopf $G-$coalgebra . ☐

Corollary 4.6. *Let (A,Δ) be a multiplier Hopf $G-$coalgebra . Then for any element $a \in A_p$ there exist some elements $e, f \in A_p$ such that $ea = a = af$. If the antipode of (A,Δ) is endomorphism of A (i.e. $S_p(A_p) \subset A_{p^{-1}}$ for all $p \in G$) then such elements e, f exist for any finitely many elements $a_1, a_2, ..., a_n$ in A_p such that $ea_i = a_i = a_i f$ for any i.*

Corollary 4.7. *If (A,Δ) is a regular multiplier Hopf $G-$coalgebra then*

1. $(\varepsilon \otimes I)((1 \otimes a)\Delta_{e,p}(b)) = ab$,
2. $(I \otimes \varepsilon)(\Delta_{p,e}(a)(b \otimes 1)) = ab$

for all $p \in G$ and $a, b \in A_p$.

Proof. Since (A,Δ) is a regular multiplier Hopf $G-$coalgebra then (A_G,Δ_G) is a regular multiplier Hopf algbera . For all $a, b \in A_p$ we have

$$\begin{aligned} ab(p) &= (I \otimes \varepsilon_G)(\Delta_G(a)(b \otimes 1))(p) \\ &= (I \otimes \varepsilon)(\Delta_G(a)(b \otimes 1)(p,e)) \\ &= (I \otimes \varepsilon)(\Delta_{p,e}(b \otimes 1)).\end{aligned}$$

Similarly, we can prove part 2. ☐

Lemma 4.8. *Let (A,Δ) be a regular multiplier Hopf $G-$coalgebra . Then*

- *There exist an antihomomorphism $S' = \{S'_p : A_p \longrightarrow M(A_{p^{-1}})\}_{p\in G}$ such that for all $a \in A_1 \quad b, c \in A_p$ and $p \in G$*

  $$m_p((I \otimes S'_{p^{-1}})((c \otimes 1)\Delta'_{p,p^{-1}}(a))(1 \otimes b)) = c\varepsilon(a)b,$$

$$m_p((c\otimes 1)(S'_{p^{-1}}\otimes I)\Delta'_{p^{-1},p}(a)(1\otimes b))=c\varepsilon(a)b.$$

- *$S_p(A_p)\subseteq A_{p^{-1}}$, and $S'_p(A_p)\subseteq A_{p^{-1}}$, S_p is invertible for all $p\in G$ with inverse $S'_{p^{-1}}$ for all $p\in G$.*

The proof of the first part is completely similar to lemma (1.7) if we use the multiplier Hopf algbera (A_G,Δ'_G) instead of (A_G,Δ_G). The 2^{nd} part is follows directly form the properties of S_G and S'_G.

Lemma 4.9. *1. Let $a\in A_{pq}$, $b\in A_q$, $a_i\in A_p$, and $b_i\in A_{q^{-1}}$. Then*

$$a\otimes S_q(b)=\sum\Delta_{pq,q^{-1}}(a_i)(1\otimes b_i)\Leftrightarrow\sum a_i\otimes S'_{q^{-1}}(b_i)=(1\otimes b)\Delta_{p,q}(a).$$

2. Let $a\in A_{pq}$, $b\in A_p$, $a_i\in A_q$, and $b_i\in A_{p^{-1}}$. Then

$$S_{p^{-1}}(b)\otimes a=\sum(b_i\otimes 1)\Delta_{p^{-1},pq}(a_i)\Leftrightarrow\Delta_{p,q}(a)(b\otimes 1)=\sum S'_{p^{-1}}(b_i)\otimes a_i.$$

3. Let $a\in A_{p^{-1}}$, $b\in A_{q^{-1}p^{-1}}$, $a_i\in A_{p^{-1}}$ and $b_i\in A_q$. Then the following statements are equivalent :

(a) $\Delta_{q,q^{-1}p^{-1}}(a)(1\otimes b)=\sum\Delta_{q,q^{-1}p^{-1}}(a_i)(b_i\otimes 1)$,

(b) $a\otimes S'_{pq}(b)=\sum(a_i\otimes 1)\Delta_{p^{-1},pq}(b_i)$,

(c) $(1\otimes a)\Delta_{q^{-1},p^{-1}}(b)=\sum S_q(b_i)\otimes a_i$.

Proof. Since (A_G,Δ_G) is a regular multiplier Hopf algbera then for all $a\in A_{pq}$, $b\in A_q$, $a_i\in A_p$, and $b_i\in A_{q^{-1}}$, we have

$$(a\otimes S_G(b))(pq,q^{-1})=(\sum\Delta_G(a_i)(1\otimes b_i))(pq,q^{-1})\Leftrightarrow(\sum a_i\otimes S_G(b_i))(p.q)=((1\otimes b)\Delta_G(a))(p,q),$$
$$a(pq)\otimes S_q(b(q))=\sum\Delta_{pq,q^{-1}}(a_i(p))(1\otimes b_i(q^{-1}))\Leftrightarrow\sum a_i(p)\otimes S'_{q^{-1}}(b_i(q^{-1}))=(1\otimes b(q))\Delta_{p,q}(a(pq)).$$

Similarly, we can prove parts 2 and 3. ☐

Lemma 4.10. *Let $b\in A_p$ and $a\in A_{pq}$. Then*

$$(1\otimes S_p(b))\Delta_{q^{-1},p^{-1}}(S_{pq}(a))=(S_q\otimes S_p)(\Delta'_{p,q}(a)(1\otimes b)).$$

Proof. Let $b\in A_p$ and $a\in A_{pq}$. Then

$$\begin{aligned}(1\otimes S_G(b))\Delta_G(S_G(a))(q^{-1},p^{-1}) &= (S_G\otimes S_G)(\Delta'_G(a)(1\otimes b))(q^{-1},p^{-1}),\\ (1\otimes S_p(b(p)))\Delta_{q^{-1},p^{-1}}(S_{pq}(a(pq))) &= \sigma((S_p\otimes S_q)(\Delta_{p,q}(a(pq))(b(p)\otimes 1)))\\ &= (S_q\otimes S_p)(\Delta'_{p,q}(a(pq))(1\otimes b(p))).\end{aligned}$$

☐

Proposition 4.11. *If (A, Δ) is a multiplier Hopf $G-$coalgebra. Define the vector space $A^\star$ by*

$$A^\star = \oplus_{\phi \in G} A^\star_\phi,$$

where A^*_ϕ is the vector space of linear functionals on A_ϕ spanned by elements of the form $a \to f(bac)$ with $b, c \in A_\phi$ and $f \in A'_\phi$. A^* can be made into associative algebra under the product

$$(fg)(a) = (f \otimes g)(\Delta_{\phi,q}(a))$$

where $f \in A^*_\phi$ and $g \in A^*_q$.

Proof. Let $f \in A^*_\phi$ and $g \in A^*_q$. Then there exist $b, c \in A_\phi$, $d, e \in A_q$, $f^{'} \in A^{'}_\phi$, and $g^{'} \in A^{'}_q$ such that $f(a) = f'(bac)$ for all $a \in A_\phi$, and $g(a) = g^{'}(dae)$ for all $a \in A_q$. Assume that $a \in A_{\phi,q}$ then

$$\begin{aligned} (fg)(a) &= m_c(f \otimes g)(\Delta_{\phi,q}(a)) \\ &= (f' \otimes g')((b \otimes d)\Delta_{\phi,q}(a)(c \otimes e)). \end{aligned}$$

Since $(b \otimes d)\Delta_{\phi,q}(a)(c \otimes e) \in A_\phi \otimes A_q$ then $(fg)(a) \in \mathbb{C}$. It means that fg is a linear functional on $A_{\phi q}$. By linearity it can be extended to all $f, g \in A^*$.

By the bijectivity of $T^1_{\phi,q}$ and $T^2_{\phi,q}$ we can write

$$\begin{aligned} b \otimes d &= \sum_{i=1}^{n} (p_i \otimes 1)\Delta_{\phi,q}(q_i), \\ c \otimes e &= \sum_{j=1}^{m} \Delta_{\phi,q}(r_j)(1 \otimes k_j). \end{aligned}$$

Then

$$(fg)(a) = \sum_{i,j} (f' \otimes g')((p_i \otimes 1)\Delta_{\phi,q}(q_i a r_j)(1 \otimes k_j)).$$

Define

$$h_{i,j} : A_{\phi q} \to \mathbb{C}$$

by

$$h_{i,j}(a) = (f' \otimes g')((p_i \otimes 1)\Delta_{\phi,q}(a)(1 \otimes k_j)).$$

Then

$$(fg)(a) = \sum_{i,j} h_{i,j}(qar).$$

i.e. $fg \in A^* \quad \forall f, g \in A^\star$ and thus $(A^*, .)$ be an algebra.

Let $f_1 \in A^*_\phi, f_2 \in A^*_q$ and $f_3 \in A^*_r$ defined by $f_i = f_i(b_i a_i c_i)$ $i = 1, 2, 3,$ where $a_1, b_1, c_1 \in A_\phi, a_2, b_2, c_2 \in A_q$ and $a_3, b_3, c_3 \in A_r$. Then

$$\begin{aligned}((f_1f_2)f_3)(a) &= (f_1f_2\otimes f_3)((1\otimes b_3)\Delta_{\phi q,r}(a)(1\otimes c_3))\\ &= (f_1\otimes f_2\otimes f_3)((b_1\otimes b_2\otimes 1)((1\otimes b_3)\Delta_{\phi q,r}(a)(1\otimes c_3))(c_1\otimes c_2\otimes 1)\\ &= (f_1\otimes f_2\otimes f_3)((1\otimes b_2\otimes b_3)((b_1\otimes 1\otimes 1)(\Delta_{\phi,q}\otimes I)(\Delta_{\phi q,r}(a)(1\otimes c_3)))\\ &\quad (c_1\otimes c_2\otimes 1))\\ &= (f_1\otimes f_2\otimes f_3)((1\otimes b_2\otimes b_3)((I\otimes \Delta_{q,\phi})((b_1\otimes 1)(\Delta_{\phi,qr}(a))(1\otimes 1\otimes c_3))\\ &\quad (c_1\otimes c_2\otimes 1))\\ &= (f_1\otimes f_2\otimes f_3)((1\otimes b_2\otimes b_3)((I\otimes \Delta_{q,\phi})((b_1\otimes 1)(\Delta_{\phi,qr}(a))(c_1\otimes 1))\\ &\quad (1\otimes c_2\otimes c_3))\\ &= (f_1(f_2f_3))(a).\end{aligned}$$

Then A^* becomes an associative algebra. □

Remark 3. By direct calculations we can prove that the two algebras $(A^*, *)$ and $({A^*}_G, m)$ are the same, where the latter is the dual algebra of (A_G, Δ_G).

4.2 Invariant functionals on regular multiplier Hopf Group coalgebras

Notation Let (A, Δ) be a regular multiplier Hopf algbera , $a \in A_{pq}$, $b \in A_q$, and $\omega \in A'_p$. Then we can define a multiplier $n \in M(A_q)$ by

$$nb = (\omega \otimes I)(\Delta_{p,q}(a)(1 \otimes b))$$
$$bn = (\omega \otimes I)((1 \otimes b)\Delta_{p,q}(a)).$$

We will write

$$n = (\omega \otimes I)\Delta_{p,q}(a).$$

Similarly, we can define $(I \otimes \omega')\Delta_{p,q}(a) \in M(A_p)$,where $\omega' \in A'_q$.

Definition 4.12. A family $\varphi = \{\varphi_p : A_p \to \mathbb{C}\}_{p\in G}$ of linear functionals on A is called left invariant if

$$(I \otimes \varphi_q)\Delta_{p,q}(a) = \varphi_{pq}(a)1_p \quad \forall a \in A_{pq}, \ \ p, q \in G.$$

A family $\varphi = \{\varphi_p\}_{p\in G}$ is said to be nonzero if $\varphi_p \neq 0$ for some $p \in G$.
A family $\psi = \{\psi_p : A_p \to \mathbb{C}\}_{p\in G}$ of linear functionals on A is called right invariant if

$$(\psi_p \otimes 1)\Delta_{p,q}(a) = \psi_{pq}(a)1_q \quad \forall a \in A_{pq}, p, q \in G$$

A family $\psi = \{\psi_p\}_{p\in G}$ is said to be nonzero if $\psi_p \neq 0$ for some $p \in G$.

Lemma 4.13. *If φ is a nonzero left invariant family of functionals then $\varphi_p \neq 0$ for all $p \in G$ such that $A_p \neq 0$.*

Proof. If $\varphi = \{\varphi_p\}_{p\in G}$ is a nonzero left invariant family of functionals then $\varphi_q \neq 0$ for some $q \in G$. Let $p \in G$ such that $A_p \neq 0$. Then $A_{qp^{-1}} \neq 0$ and we have for all $b \in A_{qp^{-1}}$

$$(I \otimes \varphi_p)\Delta_{qp^{-1},p}(a)(b \otimes 1) = \varphi_q(a)b.$$

Hence $\varphi_p \neq 0$.
Similarly, the right case can be proven. ☐

Proposition 4.14. *A has a left invariant family of functionals iff A_G has a left invariant functional.*

Proof. Let $\varphi = \{\varphi_p\}_{p\in G}$ be a left invariant family of functionals for A. Define a linear functional $\varphi_G : A_G \to \mathbb{C}$ by

$$\varphi_G(a) = \sum_{p\in G} \varphi_p(a(p)) \qquad \forall a \in A_G.$$

For $a, b \in A_G$, we have

$$\begin{aligned} [(I \otimes \varphi_G)(\Delta_G(a)(b \otimes 1))](p) &= \sum_{q\in G}(I \otimes \varphi_q)[(\Delta_G(a)(b \otimes 1))(p, q)] \\ &= \sum_{q\in G}(I \otimes \varphi_q)[\Delta_{p,q}(a(pq))(b(p) \otimes 1)] \\ &= \sum_{q\in G} \varphi_{pq}(a(pq))b(p) \\ &= (\varphi_G(a)b)(p). \end{aligned}$$

Similarly, we can prove $(I \otimes \varphi_G)((b \otimes 1)\Delta_G(a)) = b\varphi_G(a)$. Thus φ_G is left invariant.

Let φ_G is a left invariant functional for A_G. Define $\varphi_p : A_p \to \mathbb{C}$ by $\varphi_p(a(p)) = \varphi_G(a)$ for all $a \in A_p$. Let $a \in A_{pq}$ and $b \in A_p$. Then

$$\begin{aligned} (I \otimes \varphi_q)\Delta_{p,q}(a(pq))(b(p) \otimes 1) &= (I \otimes \varphi_q)([\Delta_G(a)(b \otimes 1)](p, q)) \\ &= \sum_{r\in G}(I \otimes \varphi_r)([\Delta_G(a)(b \otimes 1)](p, r)) \\ &= [(I \otimes \varphi_G)(\Delta_G(a)(b \otimes 1))](p) \\ &= \varphi_G(a)b(q) = \varphi_{pq}(a)b(q). \end{aligned}$$

Note $(I \otimes \varphi_r)([\Delta_G(a)(b \otimes 1)](p, r)) = 0 \ \ \forall r \neq q$.
Similarly, we can prove that $(I \otimes \varphi_p)((b \otimes 1)\Delta_{p,q}(a)) = b\varphi_{pq}(a)$. Thus φ is left invariant.
Similarly, the right case can be proven. ☐

Corollary 4.15. *Let (A, Δ) be a regular multiplier Hopf $G-$coalgebra with left invariant family of functionals. Then for any finitely many elements $a_q, a_2, ..., a_n \in A_p$ there exist an element $e \in A_p$ such that such that $ea_i = a_i = a_i e$ for any i*

Corollary 4.16. *If φ is left invariant then $\varphi \circ S = \{\varphi_{p^{-1}} \circ S_p : A_P \to \mathbb{C}\}_{p \in G}$ is right invariant.*

Proof. Let $\varphi = \{\varphi_p\}_{p \in G}$ be a left invariant family of functionals for A. Then φ_G is a left invariant functional for A_G. Hence, $\varphi_G \circ S_G$ is a right invariant functional for A_G. Then the family $\{\varphi_{p^{-1}} \circ S_p : A_p \to \mathbb{C}\}_{p \in G}$ is right invariant. ☐

Corollary 4.17. *If $\varphi = \{\varphi_p\}_{p \in G}$ is a nonzero left invariant family of functionals for A then φ is unique (up to scaler). The members of φ are faithful. Also, if $\psi = \{\psi_p\}_{p \in G}$ is a nonzero right invariant family of functionals for A then ψ is unique (up to scaler). The members of ψ are faithful.*

The proof of uniqueness is an immediate consequence of the uniqueness of invariant functionals on A_G. If $a \in A_p$ such that $\varphi_p(a(p)b(p)) = 0 \ \forall b \in A_p$ then $\varphi_G(ax) = 0 \ \forall \, x \in A_G$ and hence $a = 0$ (φ_G is faithful). Similarly, If $b \in A_p$ such that $\varphi_p(a(p)b(p)) = 0 \ \forall \ a \in A_p$ then $b = 0$.
If A_p is unital for all $p \in G$ then A is a Hopf $G-$coalgebra and the above corollary gives the uniqueness of its left integrals.

Let (A, Δ) be multiplier Hopf $G-$coalgebra with a left invariant family of functionals $\varphi = \{\varphi_p\}_{p \in G}$. Then (A_G, Δ_G) has a left invariant functional φ_G. Hence, there exist a multiplier $\mathcal{D}_G \in M(A_G) = \{a : G \to \cup M(A_p) \mid a(p) \in M(A_p)\}$ such that

$$(\varphi_G \otimes I)\Delta_G(a) = \varphi_G(a)\mathcal{D}_G.$$

Let $a \in A_{pq}$ and $b \in A_q$. Then

$$[(\varphi_G \otimes I)(\Delta_G(a)(1 \otimes b))](q) = \varphi_G(a)\mathcal{D}_G(q)b(q).$$

By the definition of φ_G we get

$$\begin{aligned} \varphi_{pq}(a(pq))\mathcal{D}_G(q)b(q) &= \sum_{r \in G} (\varphi_r \otimes I)(\Delta_{r,q}(a(rq))(1 \otimes b(q))) \\ &= (\varphi_p \otimes I)(\Delta_{p,q}(a(pq))(1 \otimes b(q))). \end{aligned}$$

We define a family of multipliers $\mathcal{D} = \{\mathcal{D}_p \in M(A_p)\}_{p \in G}$ by $\mathcal{D}_p = \mathcal{D}_G(p)$.

Lemma 4.18. *For all $p \in G$ there exist a multiplier $\mathcal{D}_p \in M(A_p)$ such that for all $p, q \in G$*

$$(\varphi_p \otimes I)\Delta_{p,q}(a) = \varphi_{pq}(a)\mathcal{D}_q.$$

Also, $\mathcal{D}_p$ is invertible and satisfies

$$\Delta_{p,q}(\mathcal{D}_{pq}) = \mathcal{D}_p \otimes \mathcal{D}_q, \qquad \varepsilon(\mathcal{D}_1) = 1, \qquad S_p(\mathcal{D}_p) = \mathcal{D}_{p^{-1}}^{-1}.$$

Corollary 4.19. *Let $\varphi = \{\varphi_p\}_{p\in G}$ be a nonzero left invariant family of functionals for A. Then*

1. *For all $p \in G$, $a \in A_p$*
$$\varphi_{p^{-1}}(S_p(a)) = \varphi_p(a\mathcal{D}_p).$$
2. *For all $a \in A_p$ there exist $b \in A_p$ such that*
$$\varphi_p(ca) = \varphi_p(bc) \qquad \forall c \in A_p.$$

Remark 4. Let $\varphi = \{\varphi_p\}_{p\in G}$ be a nonzero left invariant family of functionals for A. Then

1. If $\psi = \varphi \circ S$, and if we replace a by ca, $c \in A_p$ in part (1) of corollary (4.19) , we have $\psi_p(ca) = \varphi_p(ca\mathcal{D}_p)$, with $b = a\mathcal{D}_p$ we get $\psi_p(ca) = \varphi_p(cb)$.
2. In part (2) of corollary (4.19) if we apply S we get $\psi_p(ca) = \psi_p(bc)$ for all $c \in A_p$.
3. By (1),(2) we get that for all $a \in A_p$ there exist $b \in A_p$ such that $\varphi_p(ac) = \varphi_p(cb)$ for all $c \in A_p$.
4. If we combine all of this we get the equality of the four following set of functionals:
$$\{\varphi_p(a.) \mid a \in A_p\}, \qquad \{\varphi_p(.a) \mid a \in A_p\},$$
$$\{\psi_p(a.) \mid a \in A_p\}, \qquad \{\psi_p(.a) \mid a \in A_p\}.$$

Let (A, Δ) be multiplier Hopf $G-$coalgebra with a left invariant family of functionals $\varphi = \{\varphi_p\}_{p\in G}$. Then (A_G, Δ_G) has a left invariant functional φ_G. Hence, there exist an automorphism σ_G such that φ_G is σ_G invariant and $\varphi_G(ab) = \varphi_G(b\sigma_G(a))$ or all $a, b \in A_G$.
For $a, b \in A_p$ we have

$$\varphi_G(ab) = \sum_r \varphi_r(a(r)b(r)) = \varphi_p(a(p)b(p)).$$

On the other hand

$$\varphi_G(b\sigma_G(a)) = \sum_r \varphi_r(b(r)\sigma_G(a)(r)) = \varphi_p(b(p)\sigma_G(a)(p)).$$

Define $\sigma_p : A_p \longrightarrow A_p$ by $\sigma_p(a(p)) = \sigma_G(a)(p) \;\; \forall\, a \in A_p$
Note : By the faithfulness of φ_p we have that $\sigma_p(a(p)) \neq 0$ for all $0 \neq a \in A_p$.

Lemma 4.20. *There exist a family of automorphisms $\sigma = \{\sigma_p\}_{p\in G}$ of A such that $\varphi_p(ab) = \varphi_p(b\sigma_p(a))$ for all $a, b \in A_p$. We have also that φ_p is σ_p invariant.*

Corollary 4.21. *Let σ, σ' denoted the two families of automorphisms associated with φ and ψ respectively. Then for all $a \in A_{pq}$ the automorphisms σ, σ' satisfying :*

$$\begin{aligned} S_p \circ \sigma'_p &= \sigma^{-1}_{p^{-1}}, \\ \Delta_{p,q}(\sigma_{pq}(a)) &= (S_{p^{-1}} \circ S_p \otimes \sigma_{pq})\Delta_{p,q}(a), \\ \Delta_{p,q}(\sigma'_{pq}(a)) &= (\sigma'_{pq} \otimes S^{-1}_p \circ S^{-1}_{p^{-1}})\Delta_{p,q}(a). \end{aligned}$$

Corollary 4.22. *For all $a, b \in A_p$ we have*

$$\varphi_p((S_{p^{-1}} \circ S_p)(a)b) = \varphi_p(b\sigma_p((S_{p^{-1}} \circ S_p)(a))).$$

Proof. let $a, b \in A_p$. Then

$$\begin{aligned} \varphi_G(S^2(a)b) &= \varphi_p(((S \circ S)(a))(p)b(p)) \\ &= \varphi_p(((S_{p^{-1}} \circ S_p)(a(p)))b(p)). \end{aligned}$$

On the other hand, we have

$$\begin{aligned} \varphi_G(b\sigma(S^2(a))) &= \varphi_p(b(p)\sigma(((S \circ S)(a)))(p)) \\ &= \varphi_p(b(p)\sigma_p((S_{p^{-1}} \circ S_p)(a(p)))). \end{aligned}$$

Since

$$\varphi_G(S^2(a)b) = \varphi_G(b\sigma(S^2(a)))$$

we get the required formula. ☐

Let $a \in A_p$. Then by corollary (4.19) we have

$$\begin{aligned} \varphi_p(S_{p^{-1}} \circ S_p(a)) &= \varphi_{p^{-1}}(S_p(a)\mathcal{D}_{p^{-1}}) \\ &= \varphi_{p^{-1}}(S_p(\mathcal{D}^{-1}_p a)) \\ &= \varphi_{p^{-1}}(\mathcal{D}^{-1}_p a \mathcal{D}_p). \end{aligned}$$

By the uniqueness of the family φ then there exist a number τ_p such that $\varphi_p \circ S_{p^{-1}} \circ S_p = \tau_p \varphi_p$ for all $p \in G$. On the other hand $\varphi_G \circ S^2 = \sum_p \varphi_p \circ S_{p^{-1}} \circ S_p$ is left integral for A_G. Then $\varphi_G \circ S^2 = \tau_G \varphi_g$ and hence $\sum_p \tau_p \varphi_p = \sum_p \tau_G \varphi_p$ which means $\tau_p = \tau_G$ for all $p \in G$.

Corollary 4.23. *We have $\sigma_P(\mathcal{D}_P) = \sigma'_p(\mathcal{D}_p) = (1/\tau)\mathcal{D}_p$ and $\mathcal{D}_p\sigma(a) = \sigma'_p(a)$ for all $a \in A_p, p \in G$.*

Lemma 4.24. *Let (A, Δ) be a regular multiplier Hopf $G-$coalgebra , with nonzero left invariant family of functionals $\varphi = \{\varphi_p\}_{p \in G}$. Then the vector space*

$$\hat{A} = \oplus_{p \in G} \hat{A}_p,$$

where $\hat{A}_p = \{\varphi_p(.a) \mid a \in A_p\}$, can be made an associative algebra under the product

$$fg(a) = (f \otimes g)\Delta_{p,q}(a)$$

for all $f \in \hat{A}_p, g \in \hat{A}_q$, and $a \in A_{p,q}$. Moreover, the comultiplication $\hat{\Delta}$ given by

$$\begin{aligned} ((f \otimes 1)\hat{\Delta}(g))(a \otimes b) &= (f \otimes g)(\Delta_{p,q}(a)(1 \otimes b)), \\ (\hat{\Delta}(f)(1 \otimes g))(a \otimes b) &= (f \otimes g)((a \otimes 1)\Delta_{p,q}(b)), \end{aligned}$$

turn it into a regular multiplier Hopf algebra with nontrivial left and right invariant functionals given by

$$\begin{aligned} &\hat{\varphi}(f) = \varepsilon(a), \quad \textit{when} \quad f = \psi_\pi(a.) \\ &\hat{\psi}(f) = \varepsilon(a), \quad \textit{when} \quad f = \varphi_\pi(.a)\ . \end{aligned}$$

The proof is direct because the multiplier Hopf algebras $(\hat{A}, \hat{\Delta})$ and $(\hat{A}_G, \hat{\Delta}_G)$ are the same

Corollary 4.25. *Let (A, Δ) be a multiplier Hopf $G-$coalgebra with non zero left invariant family of functionals. Then the regular multiplier Hopf $G-$coalgebra associated with $(\hat{\hat{A}}, \hat{\hat{\Delta}})$ is isomorphic to (A, Δ)*

The proof is direct since (A_G, Δ_G) is isomorphic to $(\hat{\hat{A}}_G, \hat{\hat{\Delta}}_G) \cong (\hat{\hat{A}}, \hat{\hat{\Delta}})$ and combine this with Proposition (1.5).

4.3 Multiplier Hopf group coalgebras modules

Definition 4.26. Let A be a regular multiplier Hopf $G-$coalgebra. A left A-module is a family of vector spaces $R = \{R_p\}_{p\in G}$ endowed with a family of linear maps $\mu = \{\mu_p : A_p \otimes R_p \to R_p\}_{p\in G}$ such that

$$\mu_p(aa' \otimes x) = \mu_p(a \otimes \mu_p(a' \otimes x))$$

for all $a, a' \in A_p$, $x \in R_p$. We call R unital if $A_pR_p = R_p$ for all $p \in G$.

Lemma 4.27. *If R is a unital left $A-$module then $R_G = \{r : G \to \cup_{p\in G}R_P \mid r(p) \in R_p \quad \forall p \in G\}$ is a unital left A_G-module.*

Proof. Let $a \in A_G$, $r \in R_G$. We define the action of A_G on R_G by

$$(ar)(p) = a(p)r(p) \qquad \forall p \in G.$$

For $a, a' \in A_G, r \in R_G$, we have

$$((aa')r)(p) = (a(p)a'(p))r(p) = a(p)(a'(p)r(p)) = a(p)(a'r)(p) = (a(ar))(p).$$

Since $A_pR_p = R_p$ then $A_GR_G = R_G$ and hence R_G is a nuital left A_G-module. ☐

Let $\widetilde{R}$ be a unital left $\tilde{A}-$module. Then we can extend the action of $\tilde{A}$ to $M(\tilde{A})$ and hence we can define an action of $K(G)$ on $\widetilde{R}$ by

$$\delta_p r = \gamma(\delta_p)r.$$

Let $a \in \tilde{A}, r \in \widetilde{R}$ and $\delta_p, \delta_q \in K(G)$ then

$$(\delta_p\delta_q)(a.r) = \gamma(\delta_p)\gamma(\delta_q)(a.r) = \gamma(\delta_p\delta_p)(a.r) = \left\{\begin{smallmatrix} 0 & p\neq q \\ (\gamma(\delta_p)a).r); & p=q \end{smallmatrix}\right. .$$

On the other hand

$$\delta_p(\delta_q(a.r)) = \delta_p(\gamma(\delta_q)(a.r)) = \gamma(\delta_p)\gamma(\delta_q)(a.r) = \left\{\begin{smallmatrix} 0 & p\neq q \\ (\gamma(\delta_p)a).r & p=q \end{smallmatrix}\right.$$

then $(\delta_p\delta_q).r = \delta_p(\delta_q.r)$.
Moreover, since $\widetilde{R}$ is unital $\tilde{A}-$module then $\tilde{A}\widetilde{R} = \widetilde{R}$ and hence

$$K(G).\widetilde{R} = \gamma(K(G)).\tilde{A}\widetilde{R} = \tilde{A}\widetilde{R} = \widetilde{R}.$$

Thus $\widetilde{R}$ is a unital $K(G)-$module and then R can be identified with the direct of the subspaces $\{R_p = \delta_p.R\}_{p\in G}$. Hence, we get $a\gamma(\delta_p).r \in \widetilde{R}_p \ \forall r \in \widetilde{R}$

Lemma 4.28. *Let $\widetilde{R}$ be a unital left $\tilde{A}-$module. Then $\widetilde{R}$ comes from a unital left $A-$module.*

Proof. Let $\widetilde{R}$ be a unital left $\tilde{A}-$module. Define $R = \{R_p\}_{p\in G}$, where $R_p = \gamma(\delta_p)R$ and

$$\mu_p : A_p \otimes A_p \longrightarrow R_p$$

by

$$\mu_p(a\gamma(\delta_p)\gamma(\delta_p)r) = \tilde{\mu}(a\gamma(\delta_p)\gamma(\delta_p)r)$$

For $a\gamma(\delta_p), b\gamma(\delta_p) \in A_p$, $\gamma(\delta_p).r \in R_p$ we have

$$\begin{aligned} \mu_p(a\gamma(\delta_p)b\gamma(\delta_p) \otimes \gamma(\delta_p).r) &= \tilde{\mu}(a\gamma(\delta_p)b\gamma(\delta_p) \otimes \gamma(\delta_p).r) \\ &= \tilde{\mu}(a\gamma(\delta_p) \otimes \tilde{\mu}(b\gamma(\delta_p) \otimes \gamma(\delta_p).r)) \\ &= \tilde{\mu}(a\gamma(\delta_p) \otimes \mu_p(b\gamma(\delta_p) \otimes \gamma(\delta_p).r)) \\ &= \mu_p(a\gamma(\delta_p) \otimes \mu_p(b\gamma(\delta_p) \otimes \gamma(\delta_p).r)) \end{aligned}$$

Also, we have $A_pR_p = \tilde{A}\gamma(\delta_p).\gamma(\delta_p)\widetilde{R} = \gamma(\delta_p)\tilde{A}\widetilde{R} = \gamma(\delta_p)\widetilde{R} = R_p$. Hence, R is a unital left $A-$module
Define $f : \widetilde{R} \to R_G$ by $f(r)(p) = \gamma(\delta_p).r$ The map f is bijective moreover, it is satisfies

$$(\varphi(a)f(r))(p) = a\gamma(\delta_p).r = (a.r)(p) \ \ \forall a \in \tilde{A}$$

□

Lemma 4.29. *Let R be a unital left $A-$module. Then there is a unique extension to a left $M(A)-$module. We have $1_p x = x \ \forall x \in R_p, \ p \in G$.*

Let R, T are unital left $A-$modules. Then $R \otimes T$ can be made into a unital left $A-$module with structure map

$$a(x \otimes y) = \Delta_{p,q}(a)(x \otimes y), \ \ \forall a \in A_{pq}; \ x \in A_p; \ y \in A_q$$

since

$$aa'(x \otimes y) = \Delta_{p,q}(aa')(x \otimes y) = \Delta(a)\Delta(a')(x \otimes y) = \Delta(a)(\Delta(a')(x \otimes y)) = a(a'(x \otimes y)).$$

Because $\Delta_{p,q}(A_{pq})(A_p \otimes A_q) = A_p \otimes A_q$, we get

$$\begin{aligned} \Delta_{p,q}(A_{pq})(R_p \otimes T_q) &= \Delta_{p,q}(A_{pq})(A_p \otimes A_q)(R_p \otimes T_q) \\ &= (A_p \otimes A_q)(R_p \otimes T_q) \\ &= (R_p \otimes T_q). \end{aligned}$$

Then $R \otimes T$ is unital left $A-$module
Let (A, Δ) be a regular multiplier Hopf $G-$coalgebra , R' be a $G-$graded algebra over $\mathbb{C}$ with or without identity but with a nondegenerate product. Assume that $R = \{R_p\}_{p \in G}$, where $R' = \oplus_{p \in G} R_p$ is a left $A-$module.

Definition 4.30. We say that R is a left $A-$module algebra if

$$a(xx') = \sum (a_{(1,p)x})(a_{(2,q)x'}) \ \ \forall a \in A, \ x \in R_p \ , x' \in R_q.$$

Lemma 4.31. *If R is a unital left $A-$module algebra then R_G is a unital left A_G-module algebra.*

Proof. Let $a \in A_G$ and $x, x' \in R_G$ then

$$\begin{aligned} [(a_1 x)(a_2 x')] &= \sum_r (a_1 x)_r (a_2 x')_{r^{-1}p} \\ &= \sum_r (a_1(r) x(r))(a_2(r^{-1}p) x'(r^{-1}p)) \\ &= \sum_r (a_{(1,r)}(p) x(r))(a_{(2,r^{-1}p)}(p) x'(r^{-1}p)) \\ &= \sum_r a(p)(x(r) x'(r^{-1}p)) \\ &= a(p)(xx')(p) \\ &= [a(xx')](p). \end{aligned}$$

□

Let $\widetilde{R}$ be a unital left $A-$module algebra. Then $\widetilde{R}$ is a unital left $K(G)-$module algebra and hence $\widetilde{R}$ is a $G-$graded algebra.

Lemma 4.32. *Let $\widetilde{R}$ be a unital left $\tilde{A}-$module algebra. Then $\widetilde{R}$ comes from a unital $A-$module algebra.*

Proof. Let $\widetilde{R}$ be a unital left $\tilde{A}-$module. Then $a\gamma(\delta_{pq}) \in A_{pq}$, $x\gamma(\delta_p) \in A_p$ and $x'\gamma(\delta_q) \in A_q$

$$\begin{aligned} a\gamma(\delta_{pq})(\gamma(\delta_p)x\gamma(\delta_q)x') &= \sum_r (a_1\gamma(\delta_r)\gamma(\delta_p)x)(a_2\gamma(\delta_{r^{-1}pq})\gamma(\delta_q)x') \\ &= (\gamma(\delta_p)a_1x)(\gamma(\delta_q)a_2x') \\ &= m_R(\Delta_{p,q}(a\gamma(\delta_{pq}))(\gamma(\delta_p)x \otimes \gamma(\delta_q)x')). \end{aligned}$$

Then $R = \{R_p = \gamma(\delta_p)\widetilde{R}\}_{p\in G}$ is a unital left $A-$module algebra.
Moreover, the map $f : \widetilde{R} \to R_G$ defined by $f(r)(p) = \gamma(\delta_p)r$ is an algebra homomorphism since

$$f(rr')(p) = (rr')(p) = \sum_s r(s)r'(s^{-1}p) = \sum f(r)(s)f(r')(s^{-1}p) = (f(r)f(r'))(p).$$

□

Let $a \in A_G$ and $x, x' \in R_G$ then we have

$$(ax)x' = \sum a_1(x(S(a_2)x')).$$

Then for all $p \in G$, we have

$$\begin{aligned} ((ax)x')(p) &= (\sum a_1(x(S(a_2)x')))(p), \\ \sum_r (ax)(r)x'(r^{-1}p) &= \sum a_1(p)[(x(S(a_2)x'))(p)], \\ \sum_r [(a(r)x(r))x'(r^{-1}p) &= \sum a_1(p)[\sum_r x(r)[(S(a_2)x')(r^{-1}p)] \\ &= \sum_r \sum a_1(p)[x(r)[S(a_2)(p^{-1}r)x'(r^{-1}p)] \\ &= \sum_r \sum a_1(p)[x(r)[S_{p^{-1}r}(a_2(p^{-1}r))x'(r^{-1}p)]. \end{aligned}$$

Since $\Delta_G(a)(p,q) = \Delta_{p,q}(a(pq))$ then $\sum a_1(p) \otimes a_2(q) = \sum a_{(1,p)}(pq)) \otimes a_{(2,q)}(pq))$. then

$$\sum_r (a(r)x(r))x'(p^{-1}r) = \sum_r \sum a_{(1,p)}(r))[x(r)[S_{p^{-1}r}(a_{(2,p^{-1}r)}(r))x'(r^{-1}p)$$

and

$$(a(r)x(r))x'(r^{-1}p) = \sum a_{(1,p)}(r)[x(r)[S_{p^{-1}}r(a_{(2,p^{-1}r)}(r))x'(r^{-1}p)]].$$

Lemma 4.33. *1. For $a \in A_p$, $x \in R_p$ and $x' \in R_q$, we have*

$$(ax)x' = \sum a_{(1,pq)}(x(S_{q^{-1}}(a_{2,q^{-1}})x')).$$

2. For $a \in A_q$, $x \in R_p$ and $x' \in R_q$, we have

$$x(ax') = \sum a_{(2,pq)}((S_{q^{-1}}(a_{1,p^{-1}})x)x').$$

References

[1] M.E.Sweedler, Hopf algebra, Benjamin, New York, 1969.

[2] EIICHI ABE, Hopf algebras, Cambridge University Press, 1977.

[3] A.Van Daele, Multiplier Hopf algebras, Trans. Amer. Math. Soc. 342 (1994), 917-932.

[4] A.Van Daele, Discrete quantum groups, J. of Algebra 180 (1996), 431-444.

[5] V.G.Turaev, Homotopy field theory in dimention 3 and crossed group categories, preprient GT/0005291.

[6] Alexis Virelizier, Hopf group coalgebras, J. of Pure and Applied Algebra 171(2002)75-122

[7] A.S.Hegazi, A.T. Abd El-hafez and M.Mansour "Multiplier Hopf Group Coalgebra, Preprint Mansoura University 2002

[8] A.T. Abd El-hafez, L. Delvaux, and A. Van Daele, Group-cograded multiplier Hopf (*-)algebras, Preprint QA/0404026